

combination, the levels prescribed, calculated as anhydrous  $C_{10}H_{12}O_8N_2CaNa_2$ :

Food	Limitation (parts per million)	Use
Dressings, nonstandardized	75	Preservative.
French dressing .....	75	Do.
Mayonnaise .....	75	Do.
Salad dressing .....	75	Do.
Sandwich spread .....	100	Do.
Sauces .....	75	Do.

(3) Alone, as a sequestrant in the nonnutritive sweeteners that are listed in §180.37 of this chapter and that, in addition, are designed for aqueous solution: *Provided*, That the amount of the additive, calculated as anhydrous calcium disodium EDTA, does not exceed 0.1 percent by weight of the dry non-nutritive sweetener.

(c) To assure the safe use of the additive:

(1) The label and labeling of the additive container shall bear, in addition to the other information required by the act, the name of the additive.

(2) The label or labeling of the additive container shall bear adequate use directions to provide a final food product that complies with the limitations provided in paragraph (b) of this section.

(d) In the standardized foods listed in paragraphs (b) (1) and (2) of this section the additives are used only in compliance with the applicable standards of identity for such foods.

**§ 172.140 Ethoxyquin.**

(a) Ethoxyquin (1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline) may be safely used as an antioxidant for preservation of color in the production of chili powder, paprika, and ground chili at levels not in excess of 100 parts per million.

(b) In order to provide for the safe use of the additive in feed prepared in accordance with §§ 573.380 and 573.400 of this chapter, tolerances are established for residues of ethoxyquin in or on edible products of animals as follows:

- 5 parts per million in or on the uncooked fat of meat from animals except poultry.
- 3 parts per million in or on the uncooked liver and fat of poultry.
- 0.5 part per million in or on the uncooked muscle meat of animals.
- 0.5 part per million in poultry eggs.

Zero in milk.

**§ 172.145 Heptylparaben.**

(a) The food additive heptylparaben is the chemical *n*-heptyl *p*-hydroxybenzoate.

(b) It may be safely used to inhibit microbiological spoilage in accordance with the following prescribed conditions:

(1) In fermented malt beverages in amounts not to exceed 12 parts per million.

(2) In noncarbonated soft drinks and fruit-based beverages in amounts not to exceed 20 parts per million, when standards of identity established under section 401 of the Act (21 U.S.C. 341) do not preclude such use.

**§ 172.150 4-Hydroxymethyl-2,6-di-tert-butylphenol.**

The food additive 4-hydroxymethyl-2,6-di-*tert*-butylphenol may be safely used in food in accordance with the following prescribed conditions:

(a) The additive has a solidification point of 140 °C-141 °C.

(b) The additive is used as an antioxidant alone or in combination with other permitted antioxidants.

(c) The total amount of all antioxidants added to such food shall not exceed 0.02 percent of the oil or fat content of the food, including the essential (volatile) oil content of the food.

**§ 172.155 Natamycin (pimaricin).**

(a) Natamycin (CAS Reg. No. 7681-93-8), also known as pimaricin, is a polyene macrolide antimycotic substance possessing an empirical formula of  $C_{33}H_{47}NO_{13}$  and a molecular weight of 665.7.

(b) The additive shall conform to the following specifications:

Purity: 97 percent ±2 percent on an anhydrous basis.

Arsenic: Not more than 1 part per million.

Heavy metals (as Pb): Not more than 20 parts per million.

(c) The additive may be applied to the surface of cuts and slices of cheese to inhibit mold spoilage with the following limitations:

(1) The additive may be applied by dipping or by spraying, using an aqueous solution containing 200 to 300 parts per million of the additive.