

sufficient samples of urine should be taken to define the rate and extent of urinary excretion on 2 or more consecutive days to establish that steady-state conditions are achieved.

(3) A more complete characterization of the blood concentration or urinary excretion rate during the absorption and elimination phases of a single dose administered at steady-state is encouraged to permit estimation of the total area under concentration-time curves or cumulative urinary excretion-time curves and to obtain pharmacokinetic information, e.g., half-life or blood clearance, that is essential in preparing adequate labeling for the drug product.

(e) *Steady-state parameters.* (1) In certain instances, e.g., in a study involving a new drug entity, blood clearances at steady-state obtained in a multiple-dose study should be compared to blood clearances obtained in a single-dose study to support adequate dosage recommendations.

(2) In a linear system, the area under the blood concentration-time curve during a dosing interval in a multiple-dose steady-state study is directly proportional to the fraction of the dose absorbed and is equal to the corresponding "zero to infinity" area under the curve for a single-dose study. Therefore, when steady-state conditions are achieved, a comparison of blood concentrations during a dosing interval may be used to define the fraction of the active drug ingredient or therapeutic moiety absorbed.

(3) Other methods based on valid scientific reasons should be used to determine the bioavailability of a drug product having dose-dependent kinetics (non-linear system).

(f) *Measurement of an acute pharmacological effect.* When comparison of the test product and the reference material is to be based on acute pharmacological effect-time curves, measurements of this effect should be made with sufficient frequency to demonstrate a maximum effect and a lack of significant difference between the test product and the reference material.

§ 320.28 Correlation of bioavailability with an acute pharmacological effect or clinical evidence.

Correlation of in vivo bioavailability data with an acute pharmacological effect or clinical evidence of safety and effectiveness may be required if needed to establish the clinical significance of a special claim, e.g., in the case of a controlled release preparation.

§ 320.29 Analytical methods for an in vivo bioavailability study.

(a) The analytical method used in an in vivo bioavailability study to measure the concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), in body fluids or excretory products, or the method used to measure an acute pharmacological effect shall be demonstrated to be accurate and of sufficient sensitivity to measure, with appropriate precision, the actual concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), achieved in the body.

(b) When the analytical method is not sensitive enough to measure accurately the concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), in body fluids or excretory products produced by a single dose of the test product, two or more single doses may be given together to produce higher concentration if the requirements of § 320.31 are met.

§ 320.30 Inquiries regarding bioavailability and bioequivalence requirements and review of protocols by the Food and Drug Administration.

(a) The Commissioner of Food and Drugs strongly recommends that, to avoid the conduct of an improper study and unnecessary human research, any person planning to conduct a bioavailability or bioequivalence study submit the proposed protocol for the study to FDA for review prior to the initiation of the study.

(b) FDA may review a proposed protocol for a bioavailability or bioequivalence study and will offer advice with respect to whether the following conditions are met: