

B, and C of this part, of a disease condition does not imply a finding on the part of the Food and Drug Administration that any drug or device is efficacious in such condition; nor is any drug or device bearing labeling referring to such disease condition precluded from regulatory action under the applicable provisions of the act if such claim is considered to be misbranding.

§ 369.9 General warnings re accidental ingestion by children.

Section 369.20 includes under certain items, but not all medicines, the statement: "Warning—Keep this and all medicines out of children's reach. In case of accidental overdose, contact a physician immediately", or "Warning—Keep out of the reach of children". However, in view of the possibility of accidental ingestion of drugs, it is not only suggested but is recommended that one of these statements be used on the label of all drug products.

§ 369.10 Conspicuousness of warning statements.

Necessary warning statements should appear in the labeling prominently and conspicuously as compared to other words, statements, designs, and devices, and in bold type on clearly contrasting background, in order to comply with the provisions of section 502(c) and (f)(2) of the act. The warning statements should be placed in the labeling in juxtaposition with the directions for use and, in any case, should appear on the label when there is sufficient label space in addition to mandatory label information.

Subpart B—Warning and Caution Statements for Drugs

§ 369.20 Drugs; recommended warning and caution statements.

ACETANILID.

Warning—Do not exceed recommended dosage. Overdosage or continued use may result in serious blood disturbances.

ACETOPHENETIDIN CONTAINING PREPARATIONS. (See § 201.309 of this chapter.)

Warning—This medication may damage the kidneys when used in large amounts or for a long period of time. Do not take more than the recommended dosage, nor take regularly for longer than 10 days without consulting your physician.

ANESTHETICS FOR EXTERNAL USE (LOCAL ANESTHETICS). (See also § 310.201(a)(19) and (23) of this chapter.)

Caution—Do not use in the eyes. Not for prolonged use. If the condition for which this preparation is used persists or if a rash or irritation develops, discontinue use and consult physician.

ANTIHISTAMINICS FOR EXTERNAL USE (EXCEPT PREPARATIONS FOR OPHTHALMIC USE).

Caution—Do not use in the eyes. If the condition for which this preparation is used persists or if a rash or irritation develops, discontinue use and consult physician.

ANTIHISTAMINICS, ORAL. (See also § 310.201(a)(4) and (a)(24) of this chapter.)

Caution—This preparation may cause drowsiness. Do not drive or operate machinery while taking this medication. Do not give to children under 6 years of age or exceed the recommended dosage unless directed by physician.

The reference to drowsiness is not required on preparations for the promotion of sleep or on preparations that are shown not to produce drowsiness.

ANTIPERSPIRANTS.

Do not apply to broken skin. If a rash develops, discontinue use.

ANTIPYRINE.

Warning—Do not exceed recommended dosage. If skin rash appears, discontinue use and consult physician.

ANTISEPTICS FOR EXTERNAL USE.

Caution—In case of deep or puncture wounds or serious burns, consult physician. If redness, irritation, swelling, or pain persists or increases or if infection occurs discontinue use and consult physician.

The reference to wounds and burns is not required on preparations intended solely for diaper rash.

ARSENIC PREPARATIONS.