

physician, the Poison Control Center, or hospital emergency room immediately for advice.”

The following warning: Warning—Keep out of reach of children. Do not use in unconscious persons. Ordinarily, this drug should not be used if strychnine, corrosives such as alkalies (lye) and strong acids, or petroleum distillates such as kerosene, gasoline, coal oil, fuel oil, paint thinner, or cleaning fluid have been ingested.

ISOAMYLHYDROCUPREINE AND ZOLAMINE HYDROCHLORIDE RECTAL PREPARATIONS FOR EXTERNAL USE (See §310.201(a)(3) of this chapter.)

Warning—Do not use this preparation in case of rectal bleeding, as this may indicate serious disease.

NEOMYCIN SULFATE WITH A VASOCONSTRICTOR, IN NASAL PREPARATIONS (SPRAY OR DROPS).

Caution—Do not exceed recommended dosage. Do not administer to children under 3 years of age unless directed by physician.

PRAMOXINE HYDROCHLORIDE FOR EXTERNAL USE. (See §310.201(a)(19) of this chapter.)

Caution—Do not use in the eyes or nose. Not for prolonged use. Do not apply to large areas of the body. If redness, irritation, swelling, or pain persists or increases, discontinue use unless directed by a physician.

SODIUM GENTISATE. (See §§201.314, 310.201(a)(2) of this chapter.)

Warning—Do not give to children under 6 years of age or use for prolonged period unless directed by physician.

Warning—Keep this and all medications out of the reach of children; or

Warning—Keep out of the reach of children.

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

Caution—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately.

TUAMINOHEPTANE SULFATE NASAL PREPARATIONS. (See §310.201(a)(16) of this chapter.)

Caution—Do not exceed recommended dosage. Overdosage may cause nervousness, restlessness, or sleeplessness. Individuals with high blood pressure, heart disease, diabetes, or thyroid disease should use only as directed by physician. Do not use for more than 3 or 4 consecutive days unless directed by physician.

VIBESATE PREPARATIONS. (See §310.201(a)(18) of this chapter.)

Caution—Do not use but consult physician for deep or puncture wounds or serious burns. If redness, irritation, swelling, or pain persists or increases, discontinue use and consult physician.

Warning—Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° Fahrenheit may cause bursting. Never throw container into fire or incinerator.

[39 FR 11745, Mar. 29, 1974, as amended at 40 FR 8917, Mar. 3, 1975; 40 FR 13496, Mar. 27, 1975; 41 FR 10885, Mar. 15, 1976; 42 FR 22033, Apr. 29, 1977; 42 FR 36994, July 19, 1977; 44 FR 22053, Apr. 13, 1979; 44 FR 55170, Sept. 25, 1979; 52 FR 15893, Apr. 30, 1987; 52 FR 30057, Aug. 12, 1987; 52 FR 47324, Dec. 11, 1987; 55 FR 11582, Mar. 29, 1990; 57 FR 58376, Dec. 9, 1992; 59 FR 4218, Jan. 28, 1994; 61 FR 20101, May 3, 1996]

§369.22 Drugs; warning and caution statements specifically required by law.

PREPARATIONS CONTAINING HABIT-FORMING DERIVATIVES OF SUBSTANCES NAMED IN SECTION 502(d) OF THE ACT. (See §§329.1, 329.10, and 329.20 of this chapter.)

The statement “*Warning*—May be habit forming” is required to appear on the labels of all drugs containing derivatives designated in §329.1 of this chapter as habit forming, including exempt narcotic preparations described in §329.20(a) of this chapter and preparations containing one or more derivatives of barbituric acid, unless such drug is not suitable for internal use and is distributed and sold exclusively for such external use as involves no possibility of habit formation.