

(3) *Disintegration time.* Proceed as directed in § 436.212 of this chapter, using the procedure described in paragraph (e)(1) of that section.

[39 FR 18976, May 30, 1974, as amended at 49 FR 3458, Jan. 27, 1984; 50 FR 19919, May 13, 1985]

§ 440.105b Ampicillin chewable tablets.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Each ampicillin chewable tablet contains 125 milligrams or 250 milligrams of ampicillin with suitable binders, lubricants, flavorings, and colorings. Its potency is satisfactory if it is not less than 90 percent and not more than 120 percent of the number of milligrams of ampicillin that it is represented to contain. Its loss on drying is not more than 3 percent. The ampicillin used conforms to the standards prescribed by § 440.5(a)(1).

(2) *Labeling.* In addition to the labeling requirements prescribed by § 432.5 of this chapter, this drug shall be labeled “ampicillin tablets”.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) The results of tests and assays on:

(a) The ampicillin used in making the batch for potency, loss on drying, pH, ampicillin content, concordance, crystallinity, and identity.

(b) The batch for potency and loss on drying.

(ii) Samples required:

(a) The ampicillin used in making the batch: 10 packages, each containing approximately 300 milligrams.

(b) The batch: A minimum of 30 tablets.

(b) *Tests and methods of assay—(1) Potency.* Use either of the following methods; however, the results obtained from the microbiological agar diffusion assay shall be conclusive.

(i) *Microbiological agar diffusion assay.* Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: Place a representative number of tablets into a high-speed glass blender jar with sufficient 0.1M potassium phosphate buffer, pH 8.0 (solution 3), to give a stock solution of convenient concentration. Blend for 3 to 5 minutes. Remove an aliquot and fur-

ther dilute with solution 3 to the reference concentration of 0.1 microgram of ampicillin per milliliter (estimated).

(ii) *Iodometric assay.* Proceed as directed in § 436.204 of this chapter, except in paragraph (d) of that section, add 3 drops of 1.2N hydrochloric acid to both the sample and working standard solutions after the addition of 0.01N iodine solution. Prepare the sample as follows: Blend a representative number of tablets in a high-speed blender with sufficient distilled water to give a stock solution of convenient concentration. Blend for 3 to 5 minutes. Further dilute an aliquot of the stock solution with distilled water to the prescribed concentration.

(2) *Loss on drying.* Proceed as directed in § 436.200(a) of this chapter.

[39 FR 18976, May 30, 1974, as amended at 43 FR 9800, Mar. 10, 1978; 49 FR 3458, Jan. 27, 1984; 50 FR 19919, May 13, 1985]

§ 440.105c Ampicillin capsules.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Ampicillin capsules are composed of ampicillin with or without one or more buffer substances, diluents, binders, lubricants, vegetable oils, colorings, and flavorings, enclosed in a gelatin capsule. Each capsule contains 125 milligrams, 250 milligrams, or 500 milligrams of ampicillin. Its potency is satisfactory if it is not less than 90 percent and not more than 120 percent of the number of milligrams of ampicillin that it is represented to contain. The loss on drying is not more than 4.0 percent. The ampicillin used conforms to the standards prescribed by § 440.5(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The ampicillin used in making the batch for potency, loss on drying, pH, ampicillin content, concordance, crystallinity, and identity.

(b) The batch for potency and loss on drying.

(ii) Samples required: