

(6) *pH*. Proceed as directed in § 436.202 of this chapter, using the solution obtained when the product is reconstituted as directed in the labeling.

[39 FR 18976, May 30, 1974, as amended at 42 FR 59868, Nov. 22, 1977; 45 FR 22922, Apr. 4, 1980; 47 FR 22515, May 25, 1982; 50 FR 19919, May 13, 1985. Redesignated at 55 FR 277, Jan. 4, 1990]

§ 440.241b Nafcillin sodium injection.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Nafcillin sodium injection is a frozen, aqueous, iso-osmotic solution of nafcillin sodium which may contain one or more suitable and harmless buffer substances and a tonicity adjusting agent. Each milliliter contains nafcillin sodium equivalent to 20 or 40 milligrams of nafcillin. Its nafcillin content is satisfactory if it is not less than 90 percent and not more than 120 percent of the number of milligrams of nafcillin that it is represented to contain. It is sterile. It is nonpyrogenic. Its pH is not less than 6.0 and not more than 8.5. The nafcillin sodium monohydrate used conforms to the standards prescribed by § 440.41(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter. In addition, this drug shall be labeled “nafcillin sodium injection.”

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(A) The nafcillin sodium monohydrate used in making the batch for potency, moisture, pH, crystallinity, nafcillin content, and identity.

(B) The batch for nafcillin content, sterility, pyrogens, and pH.

(ii) Samples, if required by the Center for Drug Evaluation and Research:

(A) The nafcillin sodium monohydrate used in making the batch: 10 packages, each containing approximately 300 milligrams.

(B) The batch:

(1) For all tests except sterility: A minimum of 10 immediate containers.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation.

(b) *Tests and methods of assay.* Thaw the sample as directed in the labeling. The sample solution used for testing must be at room temperature.

(1) *Nafcillin content.* Proceed as directed in § 440.241a(b)(1), except use the thawed solution.

(2) *Sterility.* Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(1) of that section.

(3) *Pyrogens.* Proceed as directed in § 436.32(a) of this chapter, except inject a sufficient volume of the undiluted solution to deliver 80 milligrams of nafcillin per kilogram.

(4) *pH.* Proceed as directed in § 436.202 of this chapter, using the undiluted solution.

[55 FR 277, Jan. 4, 1990]

§ 440.249 Oxacillin sodium injectable dosage forms.

§ 440.249a Oxacillin sodium monohydrate for injection.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Oxacillin sodium monohydrate for injection is a dry mixture of oxacillin sodium monohydrate and one or more buffer substances, with or without trisodium ethylenediamine tetraacetic acid, and with or without one or more suitable and harmless preservatives. Its potency is satisfactory if it is not less than 90 percent and not more than 115 percent of the number of milligrams of oxacillin that it is represented to contain. It is sterile. It is nonpyrogenic. Its moisture content is not more than 6.0 percent. Its pH in an aqueous solution containing 30 milligrams per milliliter is not less than 6.0 and not more than 8.5. The oxacillin sodium monohydrate used conforms to the standards prescribed by § 440.49a(a)(1).

(2) *Labeling.* In addition to the labeling requirements of § 432.5 of this chapter, this drug shall be labeled “oxacillin sodium for injection”.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The oxacillin sodium monohydrate used in making the batch

§ 440.249b

21 CFR Ch. I (4–1–98 Edition)

for potency, moisture, pH, oxacillin content, crystallinity, and identity.

(b) The batch for potency, sterility, pyrogens, moisture, and pH.

(ii) Samples required:

(a) The oxacillin sodium monohydrate used in making the batch: 10 packages, each containing approximately 300 milligrams.

(b) The batch:

(1) For all tests except sterility: A minimum of 10 immediate containers.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation, or 40 immediate containers if each contains less than 600 milligrams.

(b) *Tests and methods of assay*—(1) *Potency*—(i) *Sample preparation*. Reconstitute as directed in the labeling. Then using a suitable hypodermic needle and syringe, remove all of the withdrawable contents if it is represented as a single-dose container, or, if the labeling specifies the amount of potency in a given volume of the resultant preparation, remove an accurately measured representative portion from each container. Dilute with 1 percent potassium phosphate buffer, pH 6.0 (solution 1), to give a stock solution of convenient concentration.

(ii) *Assay procedures*. Use either of the following methods; however, the results obtained from the microbiological agar diffusion assay shall be conclusive.

(a) *Microbiological agar diffusion assay*. Proceed as directed in § 436.105 of this chapter, diluting an aliquot of the stock solution with solution 1 to the reference concentration of 5 micrograms of oxacillin per milliliter (estimated).

(b) *Iodometric assay*. Proceed as directed in § 436.204 of this chapter, diluting an aliquot of the stock solution with solution 1 to the prescribed concentration.

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(1) of that section.

(3) *Pyrogens*. Proceed as directed in § 436.32(a) of this chapter, using a solution containing 20 milligrams of oxacillin per milliliter.

(4) [Reserved]

(5) *Moisture*. Proceed as directed in § 436.201 of this chapter.

(6) *pH*. Proceed as directed in § 436.202 of this chapter, using an aqueous solution containing 30 milligrams per milliliter.

[39 FR 18976, May 30, 1974, as amended at 42 FR 59868, Nov. 22, 1977; 50 FR 19918, 19919, May 13, 1985. Redesignated at 55 FR 279, Jan. 4, 1990]

§ 440.249b Oxacillin sodium injection.

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity*. Oxacillin sodium injection is a frozen aqueous, iso-osmotic solution of oxacillin sodium which may contain one or more suitable and harmless buffer substances and a tonicity adjusting agent. Each milliliter contains oxacillin sodium equivalent to 20 or 40 milligrams of oxacillin. Its oxacillin content is satisfactory if it is not less than 90 percent and not more than 115 percent of the number of milligrams of oxacillin that it is represented to contain. It is sterile. It is nonpyrogenic. Its pH is not less than 6.0 and not more than 8.5. The oxacillin sodium monohydrate used conforms to the standards prescribed by § 440.49(a)(1), except that the pH of an aqueous solution containing 30 milligrams per milliliter is not less than 4.0 and not more than 7.0.

(2) *Labeling*. It shall be labeled in accordance with the requirements of § 432.5 of this chapter. In addition, this drug shall be labeled “oxacillin sodium injection”.

(3) *Requests for certification: samples*. In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(A) The oxacillin sodium monohydrate used in making the batch for potency, moisture, pH, oxacillin content, crystallinity, and identity.

(B) The batch for oxacillin content, sterility, pyrogens, and pH.

(ii) Samples, if required by the Center for Drug Evaluation and Research:

(A) The oxacillin sodium monohydrate used in making the batch: 10 packages, each containing approximately 300 milligrams.

(B) The batch: