

§ 448.23 Cyclosporine.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Cyclosporine is a cyclic polypeptide consisting of 11 amino acids. It is a white or essentially white finely crystalline powder. It is so purified and dried that:

- (i) Its cyclosporine content is not less than 975 micrograms per milligram and not more than 1,020 micrograms per milligram on the anhydrous basis.
- (ii) Its loss on drying is not more than 3.0 percent.
- (iii) Its heavy metals content is not more than 20 parts per million.
- (iv) It passes the identity test.

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

- (i) Results of tests and assays on the batch for cyclosporine content, loss on drying, heavy metals, and identity.
- (ii) Samples, if required by the Director, Center for Drug Evaluation and Research: 10 packages, each containing approximately 500 milligrams.

(b) *Tests and methods of assay—(1) Cyclosporine content.* Proceed as directed in § 436.346 of this chapter, except prepare the working standard and sample solutions and calculate the cyclosporine content as described in paragraphs (b)(1) (i) and (ii) of this section. A typically suitable column for cyclosporine is a 250-millimeter column having an inside diameter of 4 millimeters packed with octyl silane chemically bonded to totally porous microsilica particles, 5 to 7 microns in diameter.

(i) *Preparation of working standard and sample solutions.*

NOTE: Dissolve working standards and samples immediately before analysis.

(a) *Preparation of working standard solution.* Dissolve an accurately weighed portion of the working standard in ethanol by shaking for at least 15 minutes. If necessary, ultrasonicate until the solution becomes completely clear. Dilute with ethanol to obtain a solution containing 1,000 micrograms of cyclosporine activity per milliliter.

(b) *Preparation of sample solutions.* Prepare all sample solutions as directed for preparation of working standard solutions, except dilute with ethanol to obtain a solution containing 1,000 micrograms of cyclosporine per milliliter (estimated).

(ii) *Calculations.* Calculate the micrograms of cyclosporine per milligram of sample as follows:

$$\text{Micrograms of cyclosporine per milligram} = \frac{A_u \times P_s \times 100}{A_s \times C_u \times (100 - m)}$$

where:

A_u =Area of the cyclosporine peak in the chromatogram of the sample (at a retention time equal to that observed for the standard);

A_s =Area of the cyclosporine peak in the chromatogram of the cyclosporine working standard;

P_s =Cyclosporine activity in the cyclosporine working standard solution in micrograms per milliliter;

C_u =Milligrams of cyclosporine per milliliter of sample solution; and

m =Percent loss on drying of the sample.

(2) *Loss on drying.* Proceed as directed in § 436.200(a) of this chapter.

(3) *Heavy metals.* Proceed as directed in § 436.208 of this chapter.

(4) *Identity.* The high-pressure liquid chromatogram of the sample determined as directed in paragraphs (b)(1) of this section compares qualitatively to that of the cyclosporine working standard.

[49 FR 22632, May 31, 1984, as amended at 55 FR 11584, Mar. 29, 1990]

§ 448.25 Gramicidin.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Gramicidin is the white, or nearly white, odorless, crystalline compound of a kind of gramicidin or a mixture of two or more such compounds. It is so purified and dried that:

(i) It has a potency of not less than 900 micrograms of gramicidin per milligram.

(ii) [Reserved]

(iii) Its loss on drying is not more than 3 percent.

(iv) Its residue on ignition is not more than 1.0 percent.

(v) Its melting point is not below 229° C after drying in vacuum at 60° C for 3 hours.

(vi) When calculated on the anhydrous basis, the difference between the absorptivity value at the maximum occurring at 282 nanometers and the absorptivity value at the minimum occurring at 247 nanometers is 100±4 percent of the difference obtained with the gramicidin working standard.

(vii) It is crystalline.

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5(b) of this chapter.

(3) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on the batch for potency, loss on drying, residue on ignition, melting point, identity, and crystallinity.

(ii) Samples required of the batch: Ten packages, each containing approximately 500 milligrams.

(b) *Tests and methods of assay*—(1) *Potency.* Proceed as directed in § 436.106 of this chapter, preparing the sample for assay as follows: Dissolve an accurately weighed sample in sufficient alcohol U.S.P. XX to obtain a stock solution of convenient concentration. Further dilute the stock solution volumetrically with alcohol U.S.P. XX to the reference concentration of 0.04 microgram of gramicidin per milliliter (estimated).

(2) [Reserved]

(3) *Loss on drying.* Proceed as directed in § 436.200(b) of this chapter.

(4) *Residue on ignition.* Proceed as directed in § 436.207(a) of this chapter.

(5) *Melting point.* Proceed as directed in § 436.209 of this chapter.

(6) *Identity.* Accurately weigh about 20 milligrams of the sample and dilute in ethyl alcohol to give a concentration of 0.05 milligram (estimated) of gramicidin per milliliter. Prepare a solution of the gramicidin working standard to contain 0.05 milligram per milliliter in ethyl alcohol. Using a suitable recording spectrophotometer with 1-centimeter cells, record the ultraviolet absorbance spectrum of each solution from 220 nanometers to 320 nanometers. The ultraviolet absorbance spectrum of the sample solution

should compare qualitatively to that of the working standard solution. Determine the absorptivities of each at the maximum occurring at 282 nanometers and at the minimum occurring at 247 nanometers (the exact position of the maximum and minimum of the gramicidin working standard should be determined for the particular instrument used).

(7) *Crystallinity.* Proceed as directed in § 436.203(a) of this chapter.

[39 FR 19115, May 30, 1974, as amended at 41 FR 24883, June 21, 1976; 47 FR 23710, June 1, 1982; 50 FR 19920, May 13, 1985]

§ 448.30 Polymyxin B sulfate.

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity.* Polymyxin B sulfate is the sulfate salt of a kind of polymyxin or a mixture of two or more such salts. It is a white to buff-colored powder. It is so purified and dried that:

(i) Its potency is not less than 6,000 units of polymyxin B per milligram on an anhydrous basis.

(ii) [Reserved]

(iii) Its loss on drying is not more than 7.0 percent.

(iv) Its pH is an aqueous solution containing 5 milligrams per milliliter is not less than 5.0 and not more than 7.5.

(v) It gives positive color identity tests for polymyxin.

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on the batch for potency, loss on drying, pH, and identity.

(ii) Samples required: 10 packages, each containing approximately 300 milligrams.

(b) *Tests and methods of assay*—(1) *Potency.* Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: Add 2.0 milliliters of sterile distilled water to each 5 milligrams of an accurately weighed portion of the sample. Dilute with sufficient 10 percent potassium phosphate buffer, pH 6.0 (solution 6), to give a stock solution containing 10,000 units of polymyxin B