

(5) *pH*. Proceed as directed in § 436.202 of this chapter, using the undiluted suspension.

[44 FR 1374, Jan. 5, 1979, as amended at 44 FR 30334, May 25, 1979; 50 FR 19921, May 13, 1985]

### Subpart D—Ophthalmic Dosage Forms

#### § 455.310 Chloramphenicol ophthalmic dosage forms.

##### § 455.310a Chloramphenicol ophthalmic solution.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Chloramphenicol ophthalmic solution contains in each milliliter 5 milligrams of chloramphenicol with or without one or more suitable and harmless preservatives, buffer substances, and surfactants, in an aqueous solution. Its potency is satisfactory if it is not less than 90 percent and not more than 130 percent of the number of milligrams of chloramphenicol that it is represented to contain. It is sterile. Its pH is not less than 3 nor more than 6; however, if the solution is buffered, its pH is not less than 7.0 nor more than 7.5. The chloramphenicol used conforms to the standards prescribed by § 455.10(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The chloramphenicol used in making the batch for potency, pH, specific rotation, melting range, absorptivity, and crystallinity.

(b) The batch for potency, sterility, and pH.

(ii) Samples required:

(a) The chloramphenicol used in making the batch: 10 containers, each containing approximately 300 milligrams.

(b) The batch:

(1) For all tests except sterility: A minimum of five immediate containers.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation.

(b) *Tests and methods of assay—(1) Potency.* Proceed as directed in § 436.106 of this chapter, preparing the sample for assay as follows: Dilute an accurately measured representative portion of the sample with sufficient distilled water to obtain a stock solution of convenient concentration. Further dilute an aliquot of the stock solution with distilled water to the reference concentration of 2.5 micrograms of chloramphenicol per milliliter (estimated).

(2) *Sterility.* Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(1) of that section.

(3) *pH.* Proceed as directed in § 436.202 of this chapter, using the undiluted solution.

[39 FR 19166, May 30, 1974, as amended at 43 FR 59057, Dec. 19, 1978; 46 FR 46313, Sept. 18, 1981; 48 FR 3961, Jan. 28, 1983; 50 FR 19921, May 13, 1985]

##### § 455.310b Chloramphenicol for ophthalmic solution.

(a) *Requirements for certification—(1) Standards of identity, strength, quality, and purity.* Chloramphenicol for ophthalmic solution contains 25 milligrams of chloramphenicol with one or more suitable and harmless buffer substances. When reconstituted as directed in the labeling, its potency is not less than 90 percent and not more than 130 percent of the number of milligrams of chloramphenicol that it is represented to contain. It is sterile. Its pH is not less than 7.1 and not more than 7.5. The chloramphenicol used conforms to the standards prescribed by § 455.10(a)(1).

(2) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples.* In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The chloramphenicol used in making the batch for potency, pH, specific rotation, melting range, absorptivity, and crystallinity.

(b) The batch for potency, sterility, and pH.

(ii) Samples required:

(a) The chloramphenicol used in making the batch: 10 packages, each