

§ 520.1284 Sodium liothyronine tablets.

(a) *Specifications.* Sodium liothyronine tablets consist of tablets intended for oral administration which contain liothyronine at 60 or 120 micrograms per tablet, as the sodium salt.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is indicated in cases of hypothyroidism in dogs.

(2) It is administered orally to dogs at levels up to 12.8 micrograms per kilogram of body weight per day. Dosage should be adjusted according to the severity of the condition and the response of the patient. Dosage at the total replacement level (12.8µg per kilogram of body weight) should be considered for initiating therapy and then titrated downward for optimum maintenance effect. Twice daily administration is recommended.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 56 FR 50653, Oct. 8, 1991; 60 FR 55659, Nov. 2, 1995]

§ 520.1288 Lufenuron tablets.

(a) *Specifications—(1) Dogs.* Each tablet contains either 45, 90, 204.9, or 409.8 milligrams (mg) lufenuron.

(2) *Cats.* Each regular tablet contains either 90 or 204.9 mg lufenuron, each flavor tablet contains 135 or 270 mg lufenuron.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs—(1) Amount.* 10 milligrams of lufenuron per kilogram (4.5 milligrams per pound) of body weight.

(2) *Indications for use.* For use in dogs, 6 weeks of age and older, for the prevention and control of flea populations.

(3) *Limitations.* Administer tablet(s) after or in conjunction with a full meal to ensure adequate absorption. Administer tablet(s) once a month, preferably on same date each time. All dogs in a household should be treated to achieve maximum efficacy. Because the drug has no effect on adult fleas, the concurrent use of insecticides that kill adults may be required depending on the severity of the infestation.

(d) *Conditions of use in cats—(1) Amount.* Minimum of 13.6 mg lufenuron per pound (lb) of body weight (30 mg per kilogram). Recommended 90 mg regular tablet for cats up to 6 lb of body weight, 204.9 mg regular tablet for 7 to 15 lb, 135 mg flavor tablet for up to 10 lb, 270 mg flavor tablet for 11 to 20 lb. Cats over 15 lb (regular tablet) or over 20 lb (flavor tablet) are provided the appropriate combination of tablets.

(2) *Indications for use.* For control of flea populations.

(3) *Limitations.* For oral use in cats or kittens 6 weeks of age or older, once a month, directly or broken and mixed with wet food. Administer in conjunction with a full meal to ensure adequate absorption. Treat all cats in the household to ensure maximum benefits. Because the drug has no effect on adult fleas, the concurrent use of insecticides that kill adults may be necessary depending on the severity of the infestation.

[60 FR 362, Jan. 4, 1995, as amended at 62 FR 8371, Feb. 25, 1997; 62 FR 12085, Mar. 14, 1997; 62 FR 14301, Mar. 26, 1997]

§ 520.1289 Lufenuron suspension.

(a) *Specifications.* Each individual dose pack contains either 135 or 270 milligrams of lufenuron.

(b) *Sponsor.* See No. 058198 in § 510.600(c) of this chapter.

(c) *Conditions of use in cats—(1) Amount.* Minimum of 13.6 milligrams per pound of body weight (30 milligrams per kilogram). Recommended dose of 135 milligrams for up to 10 pounds of body weight or 270 milligrams for 11 to 20 pounds. Cats over 20 pounds are provided the appropriate combination of packs.

(2) *Indications for use.* For control of flea populations.

(3) *Limitations.* For oral use in cats 6 weeks of age or older, once a month, mixed with food. Administer in conjunction with a full meal to ensure adequate absorption. Treat all cats in the household to ensure maximum benefits. Because the drug has no effect on adult fleas, the concurrent use of insecticides that kill adults may be necessary depending on the severity of the infestation.

[60 FR 20402, Apr. 26, 1995, as amended at 62 FR 8371, Feb. 25, 1997]