

§ 520.2362 Thienium closylate tablets.

(a) *Chemical name.* (N,N-Dimethyl-N-2-phenoxyethyl-N-2'-thenylammonium)-p-chlorobenzene-sulfonate.

(b) *Specifications.* Thienium closylate tablets contain thienium closylate equivalent to 500 milligrams thienium as base in each tablet.

(c) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The tablets are administered orally to dogs as a single day treatment of canine ancylostomiasis by the removal from the intestines of the adult forms of the species *Ancylostoma caninum* and *Uncinaria stenocephala* (hookworms). Dogs weighing 10 pounds and over are administered 1 tablet as a single dose. Dogs weighing 5 to 10 pounds are administered one-half tablet twice during a single day. All dosages are given for 1 day only. The treatment should be repeated after 2 or 3 weeks.

(2) Suckling puppies or recently weaned puppies weighing less than 5 pounds should not be treated with the drug. Animals that are severely infected, exhibiting evidence of intestinal hemorrhage, debilitation, and anemia, should be given supportive treatment.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 41 FR 53477, Dec. 7, 1976; 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 520.2380 Thiabendazole oral dosage forms.**§ 520.2380a Thiabendazole top dressing and mineral protein feed block.**

(a) *Chemical name.* 2-(4-Thiazolyl)benzimidazole.

(b) *Specifications.* Conforms to N.F. XII.

(c) *Sponsor.* (1) See No. 017800 in § 510.600(c) of this chapter for the sponsor of the usage provided by paragraph (e)(1)(ii) of this section.

(2) See No. 050604 in § 510.600(c) of this chapter for the sponsor of the usages provided for by paragraph (e)(1)(ii) of this section.

(3) See No. 021930 in § 510.600(c) of this chapter for the sponsor of the usage

provided for by paragraph (e)(2) of this section.

(d) *Related tolerances.* See § 556.730 of this chapter.

(e) *Conditions of use.* It is used as follows:

(1) *Horses*—(i) *Route of administration.* In feed, as a top dressing.

(a) *Amount.* 2 grams per 100 pounds of body weight.

(b) *Indications for use.* For control of large strongyles, small strongyles, pinworms, and threadworms (including members of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus*, and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, and *Strongyloides*).

(c) *Limitations.* Add to the usual feed of horses mixed into that amount of the feed normally consumed at one feeding. Warning: Not for use in horses intended for food.

(ii) *Route of administration.* In feed.

(a) *Amount.* 2 grams per 100 pounds of body weight.

(1) *Indications for use.* For control of large and small strongyles, *Strongyloides*, and pinworms of the genera *Strongylus*, *Cyathostomum*, *Cylicobrachytus* and related genera, *Craterostomum*, *Oesophagodontus*, *Poteriostomum*, *Oxyuris*, and *Strongyloides*.

(2) *Limitations.* Administer in a single dosage mixed with the normal grain ration given at one feeding. Warning: Not for use in horses intended for food.

(b) *Amount.* 4 grams per 100 pounds of body weight.

(1) *Indications for use.* For control of ascarids of the genus *Parascaris*.

(2) *Limitations.* Administer in a single dosage mixed with the normal grain ration given at one feeding. Warning: Not for use in horses intended for food.

(2) *Cattle*—(i) *Route of administration.* In feed block.

(ii) *Amount.* 3.3 percent block consumed at the recommended level of 0.11 pound per 100 pounds of body weight per day.

(iii) *Indications for use.* For control of infections of gastrointestinal roundworms (*Trichostrongylus*, *Haemonchus*, *Ostertagia* and *Cooperia*).

(iv) *Limitations.* Administer to cattle on pasture or range accustomed to mineral protein block feeding for 3