

diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[52 FR 43059, Nov. 9, 1987]

§ 520.2481 Triamcinolone acetone tablets.

(a) *Specifications.* Each tablet contains either 0.5 milligram or 1.5 milligrams of the drug.

(b) *Sponsor.* See Nos. 000010 and 053501 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use specified in this section are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is indicated for use in dogs and cats for its anti-inflammatory activity.

(2) An initial daily dosage of 0.05 milligram per pound of body weight is usually sufficient to control symptoms, although up to 0.1 milligram per pound of body weight may be given daily if response to the smaller dose is inadequate. As soon as feasible, and in any case within 2 weeks, dosage should be reduced gradually to maintenance levels of 0.0125 to 0.025 milligram per pound of body weight per day. Therapy should be discontinued by a gradual reduction in dosage after the condition has been controlled for several days. Therapy may be initiated with a single dose of sterile triamcinolone acetone suspension veterinary in which case the tablet dosage should be administered beginning 5 to 7 days after the injection or when symptoms reappear.

(3) The labeling shall comply with the requirements of § 510.410 of this chapter.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 51 FR 26002, July 18, 1986; 53 FR 40727, Oct. 18, 1988; 62 FR 35076, June 30, 1997]

§ 520.2482 Triamcinolone acetone powder.

(a) *Specifications.* Each 15 grams of triamcinolone acetone oral powder

contains 10 milligrams of triamcinolone acetone.

(b) *Sponsor.* See No. 053501 in § 510.600(c) of this chapter.

(c) *NAS/NRC status.* The conditions of use specified in this section are NAS/NRC reviewed and found effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* (1) The drug is used as an anti-inflammatory agent for horses.

(2) It is administered at a dosage of 0.005 to 0.01 milligram triamcinolone acetone per pound of body weight twice daily, sprinkled (top-dressed) on a small portion of feed. Treatment may be initiated with a single dose of sterile triamcinolone acetone suspension USP followed after 3 or 4 days with the use of triamcinolone acetone oral powder.

(3) The labeling shall comply with the requirements of § 510.410 of this chapter.

(4) Not for use in horses intended for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 24884, June 21, 1976, as amended at 50 FR 41489, Oct. 11, 1985; 51 FR 26002, July 18, 1986]

§ 520.2520 Trichlorfon oral dosage forms.

§ 520.2520a Trichlorfon oral.

(a) *Chemical name.* Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphonate.

(b) *Sponsor.* See No. 017800 in § 510.600(c) of this chapter.

(c) *Special considerations.* This drug is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(d) *Conditions of use.* (1) It is intended for use in horses for the removal of bots (*Gasterophilus spp.*), ascarids (*Parascaris equorum*), and pinworms (*Oxyuris equi*).

(2) Mix the drug, either dry or dissolved in water, in feed and administer