

(*Fasciola hepatica*); grubs (parasitic stages) (*Hypoderma bovis*, *H. lineatum*); lice (*Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*); mites (*Psoroptes ovis* (syn. *P. communis* var. *bovis*), *Sarcoptes scabiei* var. *bovis*). It is also used to control infections of *D. viviparus* and *O. Ostertagi* for 21 days after treatment, and *H. placei*, *T. axei*, *C. punctata*, *C. oncophora*, and *O. radiatum* for 14 days after treatment.

(3) *Limitations.* For subcutaneous use only. Not for intravenous or intramuscular use. Do not treat cattle within 49 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. Do not use in other animal species because severe adverse reactions, including fatalities in dogs, may result. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[55 FR 38984, Sept. 24, 1990, as amended at 62 FR 14302, Mar. 26, 1997; 62 FR 63271, Nov. 28, 1997]

§ 522.1204 Kanamycin sulfate injection.

(a) *Specifications.* Kanamycin sulfate injection veterinary conforms to the standards of identity, strength, quality, and purity prescribed by § 444.230(a) of this chapter, except that each milliliter contains either 50 or 200 milligrams of kanamycin.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in the treatment of bacterial infections due to kanamycin sensitive organisms in dogs and cats.

(2) It is administered subcutaneously or intramuscularly at 5 milligrams per pound of body weight per day in equally divided doses at 12-hour intervals.

(3) Its label shall bear an appropriate expiration date.

(4) Restricted to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 27851, July 25, 1988]

§ 522.1222 Ketamine hydrochloride injectable dosage forms.

§ 522.1222a Ketamine hydrochloride injection.

(a) [Reserved]

(b) *Specifications.* The drug is a sterile aqueous solution and each milliliter contains: Ketamine hydrochloride equivalent to 100 milligrams ketamine base activity and 1:10,000 benzethonium chloride.

(c) *Sponsors.* See Nos. 000010, 000856, 045984, and 059130 in § 510.600(c) of this chapter.

(d) [Reserved]

(e) *Conditions of use.* (1) In cats:

(i) It is used for restraint or as the sole anesthetic agent in diagnostic or minor, brief surgical procedures that do not require skeletal muscle relaxation.

(ii) It is administered intramuscularly at a recommended dose that ranges from 5 to 15 milligrams per pound of body weight depending on the effect desired.

(2) In subhuman primates:

(i) It is used for restraint.

(ii) It is administered intramuscularly at a recommended dose that ranges from 3 to 15 milligrams per kilogram of body weight depending upon the species, general condition, and age of the subject.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 59342, Dec. 23, 1975, as amended at 53 FR 27851, July 25, 1988; 59 FR 41976, Aug. 16, 1994; 59 FR 49291, Sept. 27, 1994; 60 FR 49339, Sept. 25, 1995; 62 FR 22888, Apr. 28, 1997; 62 FR 35076, June 30, 1997]

§ 522.1222b Ketamine hydrochloride with promazine hydrochloride and aminopentamide hydrogen sulfate injection.

(a) *Chemical name.* Ketamine hydrochloride, (±), -2-(*o*-chlorophenyl)-2-(methylamino) cyclohexanone hydrochloride, with promazine hydrochloride, 10-[3-(dimethylamino) propyl] phenothiazine monohydrochloride, and aminopentamide hydrogen sulfate.

(b) *Specifications.* The drug is a sterile aqueous solution and each milliliter contains: Ketamine hydrochloride equivalent to 100 milligrams ketamine base activity, 7.5 milligrams of promazine hydrochloride, and 0.0625 milligram of aminopentamide hydrogen sulfate, with 1:10,000 benzethonium chloride.

(c) *Sponsor.* See Code No. 000856 in § 510.600(c) of this chapter.