

(3) *Dogs and cats*—(i) *Amount*. 3 to 5 milligrams per pound of body weight at 12- to 24-hour intervals.

(ii) *Indications for use*—(a) *Dogs*. Treatment of upper respiratory infections such as bronchitis, tracheobronchitis, tracheitis, laryngitis, tonsillitis, and pneumonia caused by *Staphylococci* spp., hemolytic *Streptococci* spp., and *Pasteurella multocida*.

(b) *Cats*. Treatment of upper respiratory infections when caused by *Staphylococci* spp. and hemolytic *Streptococci* spp. and for feline pneumonitis when caused by tylosin susceptible organisms.

(iii) *Limitations*. For intramuscular use only. If there is no response to therapy in 5 days, diagnosis and treatment should be reassessed. Use a 50-milligram-per-milliliter solution only. Dogs and cats receiving a dose of less than 50 milligrams (1 milliliter) should be dosed with a tuberculin syringe. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 48643, Oct. 2, 1981, as amended at 47 FR 9398, Mar. 5, 1982; 50 FR 49841, Dec. 5, 1985; 50 FR 50292, Dec. 10, 1985; 53 FR 40728, Oct. 18, 1988; 59 FR 14365, Mar. 28, 1994; 62 FR 35077, June 30, 1997]

§ 522.2640b Tylosin tartrate for injection.

(a) *Specifications*. The drug is a sterile powder containing a mixture of tylosin tartrate and sodium citrate which is reconstituted to provide 25 milligrams of tylosin activity per milliliter. Tylosin as the tartrate salt, conforms to the appropriate antibiotic standard. Tylosin contains at least 95 percent tylosin as a combination of tylosin A, tylosin B, tylosin C, and tylosin D of which at least 80 percent is tylosin A as determined by a method entitled “Determination of Factor Content in Tylosin by High Performance Liquid Chromatography,” which is incorporated by reference. Copies are available from the Dockets Management Branch (HFA-305), Food and Drug Administration, rm. 1-23, 12420 Parklawn Dr., Rockville, MD 20857, or available for inspection at the Office of the Federal Register, 800 North Capitol Street, NW., suite 700, Washington, DC 20001.

(b) *Sponsor*. See No. 000986 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.740 of this chapter.

(d) *Conditions of use*—(1) *Chickens*—(i) *Amount*. 25 milligrams per 2 pounds of body weight.

(ii) *Indications for use*. As an aid in the control and treatment of chronic respiratory disease caused by *Mycoplasma gallisepticum* sensitive to tylosin.

(iii) *Limitations*. Not for use in laying chickens producing eggs for human consumption; inject under the loose skin of the neck behind the head; if no improvement is noted within 5 days, diagnosis should be reevaluated; do not treat within 3 days of slaughter.

(2) *Turkeys*—(i) *Amount*. 6.25 to 12.5 milligrams per sinus.

(ii) *Indications for use*. As an aid in the control and treatment of infectious sinusitis caused by *Mycoplasma gallisepticum* sensitive to tylosin.

(iii) *Limitations*. Do not use in laying turkeys producing eggs for human consumption; inject 6.25 milligrams to 12.5 milligrams per sinus depending on severity of condition; treatment may be repeated in 10 days if the swelling persists; do not treat within 5 days of slaughter; may be used in conjunction with tylosin in drinking water as indicated in § 520.2640(e)(2) of this chapter.

[46 FR 48643, Oct. 2, 1981, as amended at 50 FR 49841, Dec. 5, 1985; 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994]

§ 522.2662 Xylazine hydrochloride injection.

(a) *Specifications*. Xylazine hydrochloride injection is a sterile aqueous solution containing xylazine hydrochloride equivalent to 100 milligrams of xylazine in each milliliter of solution when intended for use in horses, wild deer, and elk, and 20 milligrams of xylazine per milliliter of solution when intended for use in dogs and cats.

(b) *Sponsor*. See 000856 in § 510.600(c) of this chapter for use in horses, wild deer, and elk. See 000859 and 061651 in § 510.600(c) of this chapter for use in horses, wild deer, elk, dogs, and cats. See 061690 in § 510.600(c) of this chapter for use in horses, wild deer, elk, dogs, and cats. See 000010 in § 510.600(c) of this chapter for use in horses only.

(c) *Conditions of use.* (1) The drug is used in horses, wild deer, elk, dogs, and cats to produce sedation, as an analgesic, and a preanesthetic to local anesthesia. It may also be used in horses, dogs, and cats as a preanesthetic to general anesthesia.

(2) It is administered as follows:

(i) To horses from a solution containing 100 milligrams of xylazine per milliliter, intravenously at 0.5 milligram per pound of body weight, or intramuscularly at 1.0 milligram per pound of body weight.

(ii) To dogs and cats from a solution containing 20 milligrams of xylazine per milliliter; intravenously at 0.5 milligram per pound of body weight or intramuscularly or subcutaneously at 1.0 milligram per pound of body weight. In dogs over 50 pounds, a dosage of 0.5 mg. per pound administered intramuscularly may provide sufficient sedation and/or analgesia for most procedures.

(iii) To wild deer and elk from a solution containing 100 milligrams of xylazine (as xylazine hydrochloride) per milliliter, intramuscularly, by hand syringe or syringe dart, in the heavy muscles of the croup or shoulder as follows:

(a) Fallow deer, 2 to 4 milligrams per pound.

(b) Mule deer, sika deer, and white deer, 1 to 2 milligrams per pound.

(c) Elk, 0.25 to 0.5 milligram per pound.

(3) Not to be administered to food-producing animals.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 41 FR 24884, June 21, 1976; 41 FR 28265, July 9, 1976; 53 FR 4848, Feb. 18, 1988; 53 FR 23608, June 23, 1988; 53 FR 40728, Oct. 18, 1988; 55 FR 18724, May 4, 1990; 55 FR 32616, Aug. 10, 1990; 59 FR 14367, Mar. 28, 1994; 60 FR 33110, June 27, 1995; 60 FR 35122 and 35123, July 6, 1995; 61 FR 46548, Sept. 4, 1996; 62 FR 35077, June 30, 1997]

§ 522.2670 Yohimbine injectable.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains either 2 or 5 milligrams of yohimbine (as hydrochloride).

(b) *Sponsor.* See 061690 in § 510.600(c) of this chapter for use of 2 milligrams per milliliter solution in dogs.

(1) *Amount.* 0.05 milligram per pound (0.11 milligram per kilogram) of body weight.

(2) *Indications for use.* To reverse the effects of xylazine in dogs.

(3) *Limitations.* For intravenous use in dogs only. Not for use in food-producing animals. Safety of use in pregnant dogs or in dogs intended for breeding has not been established. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(c) *Sponsor.* See 053923 in § 510.600(c) of this chapter for use of 5 milligrams per milliliter solution in deer and elk.

(1) *Amount.* 0.2 to 0.3 milligram per kilogram of body weight.

(2) *Indications for use.* As an antagonist to xylazine sedation in free ranging or confined members of the family Cervidae (deer and elk).

(3) *Limitations.* For intravenous use only. Do not use in domestic food-producing animals. Do not use for 30 days before or during hunting season. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 8543, Feb. 16, 1993, as amended at 60 FR 57832, Nov. 22, 1995]

§ 522.2680 Zeranol.

(a) *Specifications.* Each pellet contains 12 milligrams of zeranol.

(b) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.760 of this chapter.

(d) *Conditions of use.* For use as a subcutaneous ear implant as follows:

(1) *Beef cattle*—(i) *Amount.* 36 milligrams (three 12-milligram pellets) per animal.

(ii) *Indications for use*—(A) For increased rate of weight gain and improved feed conversion in weaned beef calves, growing beef cattle, feedlot steers, and feedlot heifers.

(B) For increased rate of weight gain in suckling calves.

(iii) *Limitations.* Implant subcutaneously in ear only. Do not use in bulls intended for reproduction or in dairy animals. Do not use before 1 month of age or after weaning in heifers intended for reproduction.