

(iii) *Limitations.* For subcutaneous use only. Treatment should be repeated at 24-hour intervals, continued for 48 hours after clinical signs have disappeared, for 5 to 14 days. Do not use in animals found to be hypersensitive to the drug. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(7) *Sheep*—(i) *Amount.* 0.5 to 1.0 milligram per pound (1.1 to 2.2 milligrams per kilogram) of body weight.

(ii) *Indications for use.* For treatment of sheep respiratory disease (pneumonia) associated with *Pasteurella haemolytica* and/or *P. multocida*.

(iii) *Limitations.* For intramuscular use only. Treatment should be repeated at 24 hour intervals for a total of 3 consecutive days. Additional treatments may be given on days 4 and 5 for animals which do not show satisfactory response. Use of dosages in excess of those indicated or by unapproved routes of administration may result in illegal residues in tissues. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[53 5369, Feb. 24, 1988, as amended at 55 FR 13768, Apr. 12, 1990; 56 FR 12119, Mar. 22, 1991; 57 FR 41862, Sept. 14, 1992; 59 FR 41666, Aug. 15, 1994; 59 FR 54518, Nov. 1, 1994; 60 FR 51719, Oct. 3, 1995; 61 FR 35130, July 5, 1996; 61 FR 66583, Dec. 18, 1996]

§ 522.314 Ceftiofur hydrochloride sterile suspension.

(a) *Specifications.* Each milliliter contains ceftiofur hydrochloride equivalent to 50 milligrams of ceftiofur.

(b) *Sponsor.* See No. 000009 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.113 of this chapter.

(d) *Conditions of use.* (1) *Swine*—(i) *Amount.* 3 to 5 milligrams per kilogram (1.36 to 2.27 milligrams per pound) of body weight.

(ii) *Indications for use.* For treatment and control of swine bacterial respiratory disease (swine bacterial pneumonia) associated with *Actinobacillus (Haemophilus) pleuropneumoniae*, *Pasteurella multocida*, *Salmonella choleraesuis*, and *Streptococcus suis* Type 2.

(iii) *Limitations.* For intramuscular use only. Treatment should be repeated at 24-hour intervals for a total of 3 con-

secutive days. Do not use in animals previously found to be hypersensitive to the drug. Use of dosages in excess of those indicated or route of administration other than that recommended may result in illegal residues in tissues. Safety of ceftiofur has not been determined in breeding swine. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[61 FR 29479, June 11, 1996]

§ 522.380 Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution.

(a) [Reserved]

(b)(1) *Specifications.* Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution contains 42.5 milligrams of chloral hydrate, 8.86 milligrams of pentobarbital, and 21.2 milligrams of magnesium sulfate in each milliliter of sterile aqueous solution containing water, 33.8 percent propylene glycol, and 14.25 percent ethyl alcohol.

(2) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(3) *Conditions of use.* (i) It is used for general anesthesia and as a sedative-relaxant in cattle and horses.

(ii) For intravenous use only. The drug is administered at a dosage level of 20 to 50 milliliters per 100 pounds of body weight for general anesthesia until the desired effect is produced. Cattle usually require a lower dosage on the basis of body weight. When used as a sedative-relaxant, it is administered at a level of one-fourth to one-half of the anesthetic dosage level.

(iii) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 45 FR 16482, Mar. 14, 1980]

§ 522.390 Chloramphenicol injection.

(a) *Specifications.* Each milliliter contains 100 milligrams of chloramphenicol.

(b) *Sponsor.* See Nos. 000069 and 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* *Dogs*—(1) *Amount.* 5 to 15 milligrams per pound of