

(3) If hypersensitivity to any of the components occurs treatment with this product should be discontinued and appropriate therapy instituted. Concomitant use with other drugs known to induce ototoxicity is not recommended. This preparation should not be used in conditions where corticosteroids are contraindicated. Do not administer parenteral corticosteroids during treatment with this drug. The antibiotic sensitivity of the pathogenic organism should be determined prior to use of this preparation.

(4) For use by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975. Redesignated at 41 FR 14189, Apr. 2, 1976, and amended at 52 FR 7832, Mar. 13, 1987; 61 FR 48624, Sept. 16, 1996]

§ 524.1044c Gentamicin ophthalmic ointment.

(a) *Specifications.* Each gram of sterile ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used on dogs and cats for topical treatment of conjunctivitis caused by susceptible bacteria.

(2) Apply approximately a ½-inch strip to the affected eye 2 to 4 times a day.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14188, Apr. 2, 1976, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044d Gentamicin sulfate, betamethasone valerate ointment.

(a) *Specifications.* Each gram of ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin base and betamethasone valerate equivalent to 1 milligram of betamethasone.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used or indicated for use in dogs in the treatment of acute and chronic canine otitis externa and canine infected superficial lesions caused by bacteria sensitive to gentamicin.

(2)(i) For the treatment of acute and chronic canine otitis externa the drug

is administered by instillation of 3 to 8 drops into the ear canal twice daily for 7 days. The external ear and ear canal should be properly cleaned and dried before treatment. Remove foreign material, debris, crusted exudates, etc., with suitable nonirritating solutions. Excessive hair should be clipped from the treatment area of the external ear.

(ii) For the treatment of canine infected superficial lesions, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. A sufficient amount of the drug should be applied to cover the treatment area. The drug should be administered twice daily for 7 to 14 days.

(3) If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. Concomitant use of drugs known to induce ototoxicity should be avoided. Observe patients for signs of adrenocorticoid overdosage. The antibiotic susceptibility of the pathogenic organism should be determined prior to use of this preparation. Administration of recommended doses beyond 7 days may result in delayed wound healing. Animals treated longer than 7 days should be monitored closely.

(4) For use by or on the order of a licensed veterinarian.

[47 FR 26378, June 18, 1982, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044e Gentamicin sulfate spray.

(a) *Specification.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to 1.07 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for the treatment of pink eye in cattle (infectious bovine keratoconjunctivitis) caused by *Moraxella bovis*.

(2) One actuation of the sprayer delivers 0.7 milliliter containing 0.75 milligram gentamicin. The sprayer should be held upright 3 to 6 inches from the affected eye, with the opening directed towards the eye, and pumped once. It is advisable to treat once a day for up to 3 days.

(3) Conditions other than bacterial infections of the bovine eye and infectious keratoconjunctivitis caused by *Moraxella bovis* may produce similar signs. If conditions persists or increases, discontinue use and consult veterinarian.

[48 FR 41157, Sept. 14, 1983, as amended 52 FR 7833, Mar. 13, 1987]

§ 524.1044f Gentamicin sulfate, betamethasone valerate topical spray.

(a) *Specifications.* Each milliliter of spray contains gentamicin sulfate equivalent to 0.57 milligram of gentamicin base and betamethasone valerate equivalent to 0.284 milligram of betamethasone.

(b) *Sponsor.* See Nos. 000061 and 051259 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs in the treatment of infected superficial lesions caused by bacteria sensitive to gentamicin.

(2) For the treatment of infected superficial lesions, the lesion and adjacent area should be properly cleaned before treatment. Excessive hair should be removed. Hold bottle upright 3 to 6 inches from the lesion and depress the sprayer head twice. One actuation of the sprayer delivers 0.7 milliliter of the spray. The drug should be administered with two spray actuations 2 to 4 times daily for 7 days.

(3) If hypersensitivity to any of the components occurs, treatment should be discontinued and appropriate therapy instituted. The antibiotic susceptibility of the pathogenic organism should be determined prior to use of this preparation. Administration of recommended doses beyond 7 days may result in delayed wound healing. Animals treated longer than 7 days should be monitored closely.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 740, Jan. 7, 1985, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 10220, Mar. 6, 1997]

§ 524.1044g Gentamicin sulfate, betamethasone valerate, clotrimazole ointment.

(a) *Specifications.* Each gram (g) of ointment contains gentamicin sulfate equivalent to 3 milligrams (mg)

gentamicin base, betamethasone valerate equivalent to 1 mg betamethasone, and 10 mg clotrimazole.

(b) *Sponsor.* See 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of canine otitis externa associated with yeast (*Malassezia pachydermatis*, formerly *Pityrosporum canis*) and/or bacteria susceptible to gentamicin.

(2) For 7.5 or 15 g tube, instill 4 drops of ointment twice daily into the ear canal of dogs weighing less than 30 pounds, instill 8 drops twice daily for dogs weighing 30 pounds or more. For 215 g bottle, instill 2 drops of ointment twice daily into the ear canal of dogs weighing less than 30 pounds, instill 4 drops twice daily for dogs weighing 30 pounds or more. Therapy should continue for 7 consecutive days.

(3) The external ear should be cleaned and dried before treatment. Remove foreign material, debris, crusted exudates, etc., with suitable solutions. Excessive hair should be clipped from the treatment area. If hypersensitivity occurs, treatment should be discontinued and alternate therapy instituted.

(4) Corticosteroids administered to dogs, rabbits, and rodents during pregnancy have resulted in cleft palate in offspring. Other congenital anomalies including deformed forelegs, phocomelia, and anasarca have been reported in offspring of dogs which received corticosteroids during pregnancy. Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[58 FR 38973, July 21, 1993]

§ 524.1193 Ivermectin pour-on.

(a) *Specifications.* Each milliliter of solution contains 5 milligrams of ivermectin.

(b) *Sponsor.* See No. 050604 in § 510.600(c) of this chapter.