

and sutures from the cornea, and measurement of intraocular pressure (tonometry) when glaucoma is suspected. Local applications may also be used as an aid in the removal of foreign bodies from the nose and ear canal, as an accessory in the examination and treatment of painful otitis, in minor surgery, and prior to catheterization.

(2) It is administered as follows:

(i) For removal of sutures: Instill one to two drops 2 or 3 minutes before removal of stitches.

(ii) For removal of foreign bodies from eye, ear, and nose: For ophthalmic use, instill three to five drops in the eye prior to examination; for otic use, instill five to 10 drops in the ear; for nasal use, instill five to 10 drops in each nostril every 3 minutes for three doses.

(iii) For tonometry: Instill one to two drops immediately before measurement.

(iv) As an aid in treatment of otitis: Instill two drops into the ear every 5 minutes for three doses.

(v) For minor surgery: Instill one or more drops as required.

(vi) For catheterization: Instill two to three drops with a blunt 20-gauge needle immediately before inserting catheter.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13873, Mar. 27, 1975, as amended at 50 FR 41490, Oct. 11, 1985]

§ 524.2101 Selenium disulfide suspension.

(a) *Specifications.* The product contains 0.9-percent weight in weight (w/w) selenium disulfide (1-percent weight in volume (w/v)).

(b) *NAS/NRC status.* These conditions are NAS/NRC reviewed and found effective. NADA's for similar products for these conditions of use need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(c) *Sponsors.* See 015563, 017135, 023851, and 050604 in § 510.600(c) of this chapter.

(1) *Indications for use.* For use on dogs as a cleansing shampoo and as an agent for removing skin debris associated with dry eczema, seborrhea, and non-specific dermatoses.

(2) *Amount.* One to 2 ounces per application.

(3) *Limitations.* Use carefully around scrotum and eyes, covering scrotum with petrolatum. Allow the shampoo to remain for 5 to 15 minutes before thorough rinsing. Repeat treatment once or twice a week. If conditions persist or if rash or irritation develops, discontinue use and consult a veterinarian.

[47 FR 53351, Nov. 26, 1982, as amended at 48 FR 32762, July 19, 1983; 54 FR 36962, Sept. 6, 1989; 56 FR 9623, Mar. 7, 1991; 58 FR 41025, Aug. 2, 1993]

§ 524.2140 Squalane, pyrethrins and piperonyl butoxide.

(a) *Specifications.* The drug contains 25 percent squalane (hexamethyltetracosane), 0.05 percent pyrethrins and 0.50 percent technical piperonyl butoxide.

(b) *Sponsor.* See No. 017030 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used for the treatment of ear mites in dogs and cats.

(2) It is administered as follows: Cats and dogs 5-15 pounds body weight, 4 to 5 drops in each ear daily. Dogs 16-30 pounds body weight, 5 to 10 drops in each ear daily. Dogs 30 pounds body weight and over 10 to 15 drops in each ear daily. The recommended treatment is for 7 to 10 days with repeated treatment in 2 weeks if necessary.

§ 524.2350 Tolnaftate cream.

(a) *Specifications.* The drug contains 1 percent tolnaftate (2-naphthyl-*N*-methyl-*N*-(3-tolyl) thionocarbamate) in an anhydrous cream base.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is indicated for treatment of ringworm lesions due to *Microsporium canis* and *Microsporium gypseum* in dogs and cats.

(2) A small amount of the cream is applied to the affected areas once or twice a day for 2 to 4 weeks. The areas to be treated are first cleared of exudate and the hair clipped if the areas are not already denuded. The cream is massaged into each lesion and immediate surrounding area until the cream is no longer visible.