

(ii) Amprolium plus ethopabate in accordance with § 558.58.

(iii) Bacitracin methylene disalicylate in accordance with § 558.76.

(iv) Clopidol in accordance with § 558.175.

(v) Decoquinatate in accordance with § 558.195.

(vi) Hygromycin B in accordance with § 558.274.

(vii) Monensin in accordance with § 558.355.

(viii) Roxarsone and salinomycin in accordance with § 558.550.

(ix) Robenidine hydrochloride in accordance with § 558.515.

(x) Roxarsone in accordance with § 558.530.

(xi) Salinomycin in accordance with § 558.550.

(xii) Zoalene in accordance with § 558.680.

(xiii) Tiamulin in accordance with § 558.600.

[41 FR 10995, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.128, see the List of Sections Affected in the Finding Aids section of this volume.

**§ 558.140 Chlortetracycline and sulfamethazine.**

(a) *Approvals.* Type A medicated articles: 35 grams of chlortetracycline per pound with 7.7 percent (35 grams) of sulfamethazine to 000004 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See §§ 556.150 and 556.670 of this chapter.

(c) It is used in feed for beef cattle as follows:

(1) *Amount per head per day.* Chlortetracycline, 350 milligrams plus sulfamethazine, 350 milligrams.

(2) *Indications for use.* Aid in the maintenance of weight gains in the presence of respiratory disease such as shipping fever.

(3) *Limitations.* Feed for 28 days; withdraw 7 days prior to slaughter.

[61 FR 35954, July 9, 1996]

**§ 558.145 Chlortetracycline, procaine penicillin, and sulfamethazine.**

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline per pound, 4.4 percent (20 grams) of sulfamethazine, and procaine penicillin

equivalent in activity to 10 grams of penicillin per pound to 000004 and 046573 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline per pound, 8.8 percent of sulfamethazine, and procaine penicillin equivalent in activity to 20 grams of penicillin per pound to 000004 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(3) Type C medicated feed contains in each ton, 100 grams of chlortetracycline, 50 grams of penicillin as procaine penicillin, and 100 grams of sulfamethazine.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.670 of this chapter.

(d) *Conditions of use.* (1) It is administered to swine in a Type C feed for reduction of the incidence of cervical abscesses; treatment of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery); prevention of these diseases during times of stress; maintenance of weight gains in the presence of atrophic rhinitis; growth promotion and increased feed efficiency in swine weighing up to 75 pounds.

(2) Withdraw 15 days prior to slaughter.

[40 FR 13959, Mar. 27, 1975, as amended at 43 FR 19385, May 5, 1978; 47 FR 39814, Sept. 10, 1982; 48 FR 30615, July 5, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 56 FR 14019, Apr. 5, 1991; 61 FR 18082, Apr. 24, 1996; 62 FR 14300, Mar. 26, 1997]

**§ 558.155 Chlortetracycline, sulfathiazole, penicillin.**

(a) *Approvals.* Type A medicated articles: (1) 20 grams of chlortetracycline hydrochloride, 4.4 percent (20 grams) sulfathiazole, and procaine penicillin equivalent to 10 grams of penicillin per pound to 000004 and 000010 in § 510.600(c) of this chapter.

(2) 40 grams of chlortetracycline hydrochloride, 8.8 percent (40 grams) sulfathiazole and procaine penicillin equivalent in activity to 20 grams of penicillin per pound to Nos. 000004 and 000010 in § 510.600(c) of this chapter.

(b) *Specifications.* (1) The antibiotic substance refers to the antibiotic or feed-grade antibiotic.

(2) The antibiotic activities are expressed in terms of the appropriate antibiotic standards.

(c) *Related tolerances.* See §§ 556.150, 556.510, and 556.690 of this chapter.

(d) *Conditions of use.* It is used for swine as follows:

(1) *Amount per ton.* Chlortetracycline, 100 grams plus penicillin, 50 grams plus sulfathiazole, 100 grams.

(2) *Indications for use.* For reduction of incidence of cervical abscesses. Treatment of bacterial enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis* and vibronic dysentery). Maintenance of weight gains in the presence of atrophic rhinitis. Swine 10 pounds of body weight to 6 weeks post-weaning: Increased rate of weight gain and improved feed efficiency. Swine 6 to 16 weeks post-weaning: Increased rate of weight gain.

(3) *Limitations.* For swine raised in confinement (dry-lot) or on limited pasture. Feed as sole ration. Withdraw 7 days prior to slaughter.

(b) *Related tolerances.* See § 556.160 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* It is used as follows:

(1) *Broiler chickens*—(i) *Amount per ton.* Clopidol 113.5 grams (0.0125 percent).

(a) *Indications for use.* Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Do not feed to chickens over 16 weeks of age.

(ii) *Amount per ton.* Clopidol, 113.5 grams (0.0125 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use.* Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; improved pigmentation.

(b) *Limitations.* Do not feed to chickens over 16 weeks of age; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Amount per ton.* Clopidol, 113.5 grams (0.0125 percent) plus roxarsone, 45.4 grams (0.005 percent) plus bacitracin, 4–25 grams.

(a) *Indications for use.* Aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; improved pigmentation; increased rate of weight gain.

(b) *Limitations.* Do not feed to chickens over 16 weeks of age; withdraw 5 days before slaughter; as sole source of organic arsenic; as bacitracin methylene disalicylate, provided by No. 046573 in § 510.600(c) of this chapter; or as zinc bacitracin provided by Nos. 000061 and 046573 in § 510.600(c) of this chapter.

(iv) *Amount per ton.* Clopidol, 113.5 grams (0.0125%) plus zinc bacitracin, 5 to 25 grams.

(a) *Indications for use.* For increased rate of weight gain and improved feed efficiency; aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Feed continuously as sole ration. Zinc bacitracin as provided by Nos. 000061 and 046573 of § 510.600(c) of this chapter.

MINIMUM AMOUNT OF TYPE C FEED WHICH THE ANIMAL SHOULD CONSUME

Type of feed	Approximate body weight in pounds	Minimum desired daily feed intake in pounds
Prestarter (up to 6 weeks postweaning) .....	20	1
Starter (up to 6 weeks postweaning) .....	50	1½
Grower (6–16 weeks postweaning)	80	2
Finisher (6–16 weeks postweaning) .....	150	3

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7397, Mar. 3, 1986; 51 FR 28547, Aug. 8, 1986; 52 FR 2684, Jan. 26, 1987; 61 FR 2415, Jan. 26, 1996; 62 FR 35077, June 30, 1997; 62 FR 67725, Dec. 30, 1997; 63 FR 11599, Mar. 10, 1998]

**§ 558.175 Clopidol.**

(a) *Approvals.* Type A medicated articles: (1) 25 percent to 011526 in § 510.600(c) of this chapter.

(2) 25 percent of clopidol, 10 percent of roxarsone, and 4, 10, 15, or 25 grams of bacitracin methylene disalicylate per pound to 011526 in § 510.600(c) of this chapter.