

equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Narasin as provided by 000986, bacitracin methylene disalicylate by 046573 in § 510.600(c) of this chapter.

(vii) *Amount per ton.* Narasin 54 to 72 grams, bambermycins 1 to 2 grams, and roxarsone 22.7 to 45.4 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima*, and *E. brunetti*, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation in broiler chickens.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not feed to laying hens. Do not allow adult turkeys or horses or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdosage or lack of water intake may result in leg weakness or paralysis. Withdraw 5 days before slaughter. Narasin as provided by 000986 in § 510.600(c) of this chapter, bambermycins by 012799, and roxarsone by 046573.

(2) [Reserved]

[51 FR 29098, Aug. 14, 1986, as amended at 53 FR 24260, June 28, 1988; 54 FR 1928, Jan. 18, 1989; 54 FR 5930, Feb. 7, 1989; 54 FR 9430, Mar. 7, 1989; 54 FR 29335, July 12, 1989; 55 FR 8460, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992; 57 FR 32176, July 21, 1992; 63 FR 13124, Mar. 18, 1998]

§ 558.365 Nequinat.

(a) *Approvals.* Type A medicated articles: 4 percent to 017800 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.440 of this chapter.

(c) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Conditions of use.* It is used as follows:

(1) *Broiler or fryer chickens—(i) Amount per ton.* Nequinat, 18.16 grams.

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E.*

tenella, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Feed continuously as the sole ration.

(ii) *Amount per ton.* Nequinat, 18.16 grams (0.002 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; for improving pigmentation.

(b) *Limitations.* Feed continuously as sole ration throughout the starting period; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Amount per ton.* Nequinat, 18.16 grams (0.002 percent) plus oxytetracycline, 200 grams.

(a) *Indications for use.* For control of complicated chronic respiratory disease (air-sac infection), infectious synovitis, and treatment of blue comb (nonspecific infectious enteritis).

(b) *Limitations.* As monoalkyl (C₈–C₁₈) trimethylammonium oxytetracycline as provided by No. 000069 in § 510.600(c) of this chapter.

(2) *Roaster chickens or replacement chickens for caged layers—(i) Amount per ton.* Nequinat, 18.16 grams (0.002 percent).

(ii) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(iii) *Limitations.* Feed continuously as the sole ration; do not feed to chickens over 16 weeks of age.

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2685, Jan. 26, 1987]

§ 558.366 Nicarbazin.

(a) Type A medicated articles: 25 percent to 000986, 060728, and 063271 in § 510.600(c) of this chapter for use as indicated in the table in paragraph (c) of this section.

(b) *Related tolerances.* See § 556.445 of this chapter.

(c) *Conditions of use.* It is used in chicken feed as follows: