

equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Narasin as provided by 000986, bacitracin methylene disalicylate by 046573 in § 510.600(c) of this chapter.

(vii) *Amount per ton.* Narasin 54 to 72 grams, bambermycins 1 to 2 grams, and roxarsone 22.7 to 45.4 grams.

(A) *Indications for use.* For prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. mivati*, *E. acervulina*, *E. maxima*, and *E. brunetti*, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation in broiler chickens.

(B) *Limitations.* For broiler chickens only. Feed continuously as sole ration. Do not feed to laying hens. Do not allow adult turkeys or horses or other equines access to formulations containing narasin. Ingestion of narasin by these animals has been fatal. Use as sole source of organic arsenic. Poultry should have access to drinking water at all times. Drug overdosage or lack of water intake may result in leg weakness or paralysis. Withdraw 5 days before slaughter. Narasin as provided by 000986 in § 510.600(c) of this chapter, bambermycins by 012799, and roxarsone by 046573.

(2) [Reserved]

[51 FR 29098, Aug. 14, 1986, as amended at 53 FR 24260, June 28, 1988; 54 FR 1928, Jan. 18, 1989; 54 FR 5930, Feb. 7, 1989; 54 FR 9430, Mar. 7, 1989; 54 FR 29335, July 12, 1989; 55 FR 8460, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992; 57 FR 32176, July 21, 1992; 63 FR 13124, Mar. 18, 1998]

§ 558.365 Nequinatate.

(a) *Approvals.* Type A medicated articles: 4 percent to 017800 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.440 of this chapter.

(c) *Special considerations.* Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Conditions of use.* It is used as follows:

(1) *Broiler or fryer chickens—(i) Amount per ton.* Nequinatate, 18.16 grams.

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E.*

tenella, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(b) *Limitations.* Feed continuously as the sole ration.

(ii) *Amount per ton.* Nequinatate, 18.16 grams (0.002 percent) plus roxarsone, 45.4 grams (0.005 percent).

(a) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, and *E. mivati*; growth promotion and feed efficiency; for improving pigmentation.

(b) *Limitations.* Feed continuously as sole ration throughout the starting period; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Amount per ton.* Nequinatate, 18.16 grams (0.002 percent) plus oxytetracycline, 200 grams.

(a) *Indications for use.* For control of complicated chronic respiratory disease (air-sac infection), infectious synovitis, and treatment of blue comb (nonspecific infectious enteritis).

(b) *Limitations.* As monoalkyl (C₈–C₁₈) trimethylammonium oxytetracycline as provided by No. 000069 in § 510.600(c) of this chapter.

(2) *Roaster chickens or replacement chickens for caged layers—(i) Amount per ton.* Nequinatate, 18.16 grams (0.002 percent).

(ii) *Indications for use.* An aid in the prevention of coccidiosis caused by *E. tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

(iii) *Limitations.* Feed continuously as the sole ration; do not feed to chickens over 16 weeks of age.

[40 FR 13959, Mar. 27, 1975, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2685, Jan. 26, 1987]

§ 558.366 Nicarbazin.

(a) Type A medicated articles: 25 percent to 000986, 060728, and 063271 in § 510.600(c) of this chapter for use as indicated in the table in paragraph (c) of this section.

(b) *Related tolerances.* See § 556.445 of this chapter.

(c) *Conditions of use.* It is used in chicken feed as follows:

Nicarbazin in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
27 to 45	Narasin 27 to 45	Broiler chickens; prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. acervulina</i> , <i>E. maxima</i> , <i>E. brunetti</i> , and <i>E. mivati</i> .	Sec. 558.363 (d)(1)(iii)	000986
113.5 (0.0125 pct)	Chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis..	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; do not use as a treatment for coccidiosis; do not use in flushing mashers; do not feed to laying hens; withdraw 4 days before slaughter.	000986 060728 063271
	Bacitracin methylene disalicylate 30.	Broiler chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis; for increased rate of weight gain and improved feed efficiency.do	060728 063271
	Lincomycin 2 (0.00044 pct).	Broiler chickens; aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> and <i>E. brunetti</i>) coccidiosis; for increased rate of weight gain.do	060728 063271
	Roxarsone 22.7 (0.0025)..do	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; as sole source of organic arsenic; do not use a treatment for coccidiosis; do not use in flushing mashers; do not feed to laying hens; withdraw 5 days before slaughter.	060728 063271
	Roxarsone 22.7 (0.0025) plus lincomycin 2 (0.0004)..dodo	060728 063271

[42 FR 56729, Oct. 28, 1977; 43 FR 1942, Jan. 13, 1978, as amended at 44 FR 40887, July 13, 1979; 50 FR 13562, Apr. 5, 1985; 51 FR 7399, Mar. 3, 1986; 54 FR 1928, Jan. 18, 1989; 60 FR 29483, June 5, 1995; 61 FR 1832, Jan. 24, 1996; 61 FR 14021, Mar. 29, 1996; 61 FR 14483, Apr. 2, 1996; 62 FR 29011, May 29, 1997; 63 FR 13124, Mar. 18, 1998]

§ 558.369 Nitarsone.

(a) *Approvals.* Type A medicated articles: 50 percent to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *NAS/NRC status.* These conditions of use are NAS/NRC reviewed and found effective. NADA's for these uses may not require effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* It is used for chickens and turkeys as follows:

(1) *Amount.* Nitarsone, 0.01875 percent.

(2) *Indications for use.* As an aid in the prevention of blackhead.

(3) *Limitations.* Early medication is essential to prevent spread of disease. Adequate drinking water must be provided near feeder at all times. The drug is not effective in preventing blackhead in birds infected more than 4 or 5 days. Discontinue use 5 days before slaughtering animals for human consumption to allow elimination of the