

(1) *Indications for use.* Maintaining weight gains and feed efficiency in presence of atrophic rhinitis.

(2) *Limitations.* As tylosin phosphate.

(2) Tylosin may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Hygromycin B as in § 558.274.

(ii) Melengestrol acetate alone or in combination with certain ionophores in accordance with § 558.342.

(iii) [Reserved]

(iv) Monensin in accordance with § 558.355.

(v) Pyrantel tartrate in accordance with § 558.485.

(e) *Amount per ton.* Tylosin 100 grams. (1) *Indications for use.* Prevention and/or control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

(2) *Limitations.* As tylosin phosphate, administer for 21 days.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.625, see the List of CFR Sections Affected in the Finding Aids section of this volume.

**§ 558.630 Tylosin and sulfamethazine.**

(a) [Reserved]

(b) *Approvals.* Type A medicated article levels, a combination of equal amounts of tylosin and sulfamethazine, granted to firms as sponsor(s) and identified by drug listing numbers in § 510.600(c) of this chapter for the conditions of use indicated in paragraph (f) of this section.

(1) To 000986: 40 grams per pound each, paragraph (f)(2)(i).

(2) To 000986: 10 grams per pound each, paragraph (f)(2)(i).

(3) To 017519: 10 grams per pound each, paragraph (f)(2)(ii) of this section.

(4) To 021780: 2 grams per pound each, paragraph (f)(2)(ii) of this section.

(5) To 017800: 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(6) To 017139: 4, 10, or 20 grams per pound each, paragraph (f)(2)(ii) of this section.

(7) To 021930: 2 grams per pound each, paragraph (f)(2)(i) of this section; 5, 10, 20, or 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(8) To 017519, 026186: 5 or 10 grams per pound each, paragraph (f)(2)(ii) of this section.

(9) [Reserved]

(10) To 010439, 011749, 016968, 017473, 017519, 017790, 021780, 024174, 030841, 034936, 035098, 043727, 043733, 046573, 046987, 050568, 050639, and 051359, 053389: 5, 10, 20, or 40 grams per pound each, paragraph (f)(2)(ii) of this section.

(c)-(d) [Reserved]

(e) *Related tolerances.* See §§ 556.670 and 556.740 of this chapter.

(f) *Conditions of use.* It is used in feed for swine as follows:

(1) *Amount per ton.* Tylosin, 100 grams plus sulfamethazine, 100 grams.

(2) *Indications for use.* (i) Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica* rhinitis; prevention of swine dysentery (vibriotic); control of swine pneumonias caused by bacterial pathogens (*P. multocida* and/or *C. pyogenes*); for reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E Streptococci. Only the sulfamethazine portion of this combination is active in controlling jowl abscesses.

(ii) Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis; lowering the incidence and severity of *Bordetella bronchiseptica* rhinitis; prevention of swine dysentery (vibriotic); control of swine pneumonias caused by bacterial pathogens (*Pasteurella multocida* and/or *Corynebacterium pyogenes*).

(3) *Limitations.* As tylosin phosphate; withdraw 15 days before slaughter.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.630, see the List of CFR Sections Affected in the Finding Aids section of this volume.

**§ 558.635 Virginiamycin.**

(a) [Reserved]

(b) *Approvals.* Type A medicated articles. (1) 1.1 percent activity (5 grams per pound), 2.2 percent activity (10 grams per pound), 4.4 percent activity (20 grams per pound), 11 percent activity (50 grams per pound), and 50 percent activity (227 grams per pound) used as in paragraph (f) of this section; and 30 percent activity (136.2 grams per

pound) for the manufacture of Type C medicated feed for cattle used as in paragraph (f)(3); to 000069 in § 510.600(c) of this chapter.

(2) 2.2 percent activity (10 grams per pound) to 011490, 016968, and 017790 in § 510.600(c) of this chapter for use as in paragraphs (f)(1)(iv) and (f)(1)(v) of this section.

(c) [Reserved]

(d) *Related tolerances.* See § 556.750 of this chapter.

(e) *Special considerations.* (1) Not for use in breeding swine over 120 pounds.

(2) Dilute Type A article with at least 10 pounds of a feed ingredient prior to final mixing in 1 ton of Type C feed.

(3)-(4) [Reserved]

(f) *Conditions of use*—(1) *Swine.* It is used as follows:

(i) 100 grams per ton for 2 weeks, for treatment of swine dysentery in non-breeding swine over 120 pounds.

(ii) 100 grams per ton for 2 weeks, 50 grams per ton thereafter, for treatment and control of swine dysentery in swine up to 120 pounds.

(iii) 25 grams per ton, as an aid in control of dysentery in swine up to 120 pounds. For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred.

(iv) 10 grams per ton from weaning up to 120 pounds for increased rate of weight gain and improved feed efficiency, followed by 5 grams per ton to market weight for increased rate of weight gain and improved feed efficiency. For continuous use from weaning to market weight.

(v) 10 grams per ton from weaning up to 120 pounds for increased rate of weight gain and improved feed efficiency, followed by 5 to 10 grams per ton to market weight for increased rate of weight gain. For continuous use from weaning to market weight.

(2) *Poultry.* It is used as follows:

(i) 5 to 15 grams per ton for increased rate of weight gain, for use in broiler chickens, not for use in layers.

(ii) 5 grams per ton for increased rate of weight gain and improved feed effi-

ciency in broiler chickens, not for use in layers.

(iii) 20 grams per ton for prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to virginiamycin in broiler chickens; not for use in layers.

(iv) 10 to 20 grams per ton for increased rate of weight gain and improved feed efficiency in growing turkeys.

(3) *Cattle.* It is used as follows:

(i) 16.0 to 22.5 grams per ton to provide 100 to 340 milligrams per head per day for increased rate of weight gain.

(ii) 13.5 to 16.0 grams per ton to provide 85 to 240 milligrams per head per day for reduction of incidence of liver abscesses.

(iii) 11.0 to 16.0 grams per ton to provide 70 to 240 milligrams per head per day for improved feed efficiency.

(iv) Feed continuously as sole ration to cattle fed in confinement for slaughter. Not for use in animals intended for breeding.

(4) Virginiamycin may be used in accordance with the provisions of this section in the combinations provided, as follows:

(i) Monensin sodium in accordance with § 558.355.

(ii) Lasalocid sodium in accordance with § 558.311.

(iii) Monensin and roxarsone as in § 558.355.

(iv) Amprolium and ethopabate as in § 558.58.

(v) Halofuginone as in § 558.265.

(vi) Salinomycin alone or with roxarsone as in § 558.550.

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.635, see the List of CFR Sections Affected in the Finding Aids section of this volume.

#### § 558.680 Zoalene.

(a) *Approvals.* Type A medicated articles: 25 percent to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.770 of this chapter.

(c) *Conditions of use*—(1) *Chickens and turkeys:*