

§ 558.76

21 CFR Ch. I (4–1–98 Edition)

(iii) Bacitracin zinc in accordance with § 558.78.

(iv) Bacitracin and zoalene in accordance with § 558.680.

(v) Zoalene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986; 54 FR 18281, Apr. 28, 1989; 56 FR 19268, Apr. 26, 1991; 60 FR 39847, Aug. 4, 1995]

§ 558.76 Bacitracin methylene disalicylate.

(a) *Approvals.* Type A medicated articles: 10, 25, 30, 40, 50, 60, or 75 grams per pound to 046573 in § 510.600(c) of this chapter.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.

(d) *Conditions of use.* (1) It is used as follows:

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 4 to 50	Chickens, turkeys, and pheasants; increased rate of weight gain and improved feed efficiency ¹	046573
(ii) 5 to 20	Quail not over 5 weeks of age; increased rate of weight gain and improved feed efficiency ¹	046573
(iii) 10 to 25	Chickens; for increased egg production and improved feed efficiency for egg production.	For first 7 months of production	046573
(iv) 10 to 30	Swine: for increased rate of weight gain and improved feed efficiency.	For growing and finishing swine	000004 and 046573
	Chlortetracycline approximately 400, varying with body weight and food consumption to provide 10 milligrams per pound of body weight per day.	Swine: for increased rate of weight gain and improved feed efficiency; for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline.	Feed for not more than 14 days to provide 10 milligrams of chlortetracycline per pound of body weight per day; as chlortetracycline provided by Nos. 000004 and 046573 in § 510.600(c) of this chapter.	000004 and 046573
(v) [Reserved]				
(vi) 50	Broiler chickens; as an aid in the prevention of necrotic enteritis caused or complicated by <i>Clostridium spp.</i> or other organisms susceptible to bacitracin.	046573
(vii)—(viii) [Reserved]				
(ix) 100 to 200	Broiler chickens; as an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium spp.</i> or other organisms susceptible to bacitracin.	046573
(x) 200	Turkeys; as an aid in the control of transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.	046573
	Quail; for the prevention of ulcerative enteritis in growing quail due to <i>Clostridium colinum</i> susceptible to bacitracin methylene disalicylate	From Type A medicated articles containing 25, 40, or 50 grams of bacitracin methylene disalicylate. Feed continuously as the sole ration.	046573

Bacitracin methylene disalicylate in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(xi) 250	1. Growing/Finishing Swine: For control of swine dysentery associated with <i>Treponema hyodysenteriae</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred; or following an approved treatment of the disease condition.	As the sole ration. Not for use in swine weighing more than 250 pounds. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.	046573
	2. Pregnant sows: For control of clostridial enteritis caused by <i>C. perfringens</i> in suckling piglets..	As the sole ration. Feed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.	

¹ These conditions are NAS/NRC reviewed and found effective. Applications for these uses may not require effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(2) It is used as bacitracin methylene disalicylate in feed for animals as follows:

(i) *Amount.* 70 milligrams per head per day.

(a) *Indications for use.* Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) *Limitations.* Administer continuously throughout the feeding period.

(ii) *Amount.* 250 milligrams per head per day.

(a) *Indications for use.* Feedlot beef cattle; reduction in the number of liver condemnations due to abscesses.

(b) *Limitations.* Administer continuously for 5 days then discontinue for subsequent 25 days, repeat the pattern during the feeding period.

(3) It is used as bacitracin methylene disalicylate in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium with ethopabate in accordance with § 558.58.

(iii) Arsanilic acid with zoalene in accordance with § 558.680

(iv) Carbarson (not U.S.P.) in accordance with § 558.120.

(v) Nicarbazin as in § 558.366.

(vi) Hygromycin B in accordance with § 588.274.

(vii) Monensin in accordance with § 588.355.

(viii) Lasalocid sodium alone or with roxarsone as in § 558.311.

(ix) Monensin and roxarsone in accordance with § 558.355.

(x) Salinomycin alone or with roxarsone as in § 558.550.

(xi) Halofuginone hydrobromide and roxarsone in accordance with § 558.265.

(xii) Halofuginone in accordance with § 558.265.

(xiii) Narasin alone or in combination with roxarsone as in § 558.363.

(xiv) Semduramicin alone or in combination with roxarsone as in § 558.555.

[41 FR 10993, Mar. 15, 1976]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 558.76, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§ 558.78 Bacitracin zinc.

(a) *Approvals.* To sponsors listed in § 510.600(c) of this chapter for use as in paragraph (d) of this section as follows:

(1) To 046573: 50 grams per pound as in paragraphs (d)(1)(i), (d)(1)(ii), (d)(1)(iii), (d)(1)(iv), and (d)(2) of this section.

(2) To 000004: 10, 25, 40, and 50 grams per pound as in paragraphs (d)(1)(i), (d)(1)(ii), (d)(1)(v), (d)(1)(vi), (d)(2), and (d)(3) of this section.

(3) To 000010: 5 and 50 grams per pound for chickens as in paragraph (d)(1)(i) of this section.

(b) *Special considerations.* The quantities of antibiotics are expressed in terms of the equivalent amount of antibiotic standard.

(c) *Related tolerances.* See § 556.70 of this chapter.