

fermented ammoniated condensed whey shall not exceed 80 percent of the free-choice liquid feed.

(e) *Labeling.* The label shall bear, in addition to other information required by the act:

(1) The name of the additive.

(2) The maximum percentage of equivalent crude protein from non-protein nitrogen.

(3) Adequate directions for use in accordance with the provisions in paragraph (d) of this section.

[43 FR 33708, July 1, 1978, as amended at 46 FR 49115, Oct. 6, 1981]

§ 573.460 Formaldehyde.

The food additive formaldehyde may be safely used in the manufacture of animal feeds in accordance with the following conditions:

(a)(1) The additive is used, or intended for use, to improve the handling characteristics of animal fat in combination with certain oilseed meals by producing therefrom a dry, free-flowing product as follows:

(i) An aqueous blend of soybean and sunflower meals in a ratio of 3:1, respectively, is mixed with animal fat such that the oilseed meals and animal fat are in a ratio of 3:2. The feed ingredients are those defined by the "Official Publication" of the Association of American Feed Control Officials, Inc., 1976 ed., pages 86, 103, and 109.¹

(ii) Formaldehyde (37 percent solution) is added to the mixture at a level of 4 percent of the dry matter weight of the oilseed meals and animal fat. This mixture, upon drying, contains not more than 1 percent formaldehyde and not more than 12 percent moisture.

(2) The dried mixture described in paragraph (a) of this section is used, or intended for use, as a component of dry, nonpelleted feeds for beef and non-lactating dairy cattle.

(3) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling of the dried mixture described in paragraph (a) of this section shall bear:

(i) The name of the additive.

(ii) Adequate directions for use providing that feed as consumed is not to contain more than 25 percent of the mixture.

(b)(1) The food additive is formaldehyde (37 percent aqueous solution). It is used at the rate of 5.4 pounds (2.5 kilograms) per ton of poultry feed. At this level, it is an antimicrobial agent used to maintain complete poultry feeds salmonella negative for up to 14 days.

(2) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling shall contain:

(i) The name of the additive.

(ii) A statement that formaldehyde solution which has been stored below 40 °F or allowed to freeze should not be applied to complete poultry feeds.

(iii) Adequate directions for use including a statement that formaldehyde should be thoroughly mixed into complete poultry feed and that the finished poultry feed shall be labeled as containing formaldehyde.

(3) To assure safe use of the additive, in addition to the other information required by the Act, the label and labeling shall contain:

(i) Appropriate warnings and safety precautions concerning formaldehyde.

(ii) Statements identifying formaldehyde as a poison with potentials for adverse respiratory effects.

(iii) Information about emergency aid in case of accidental inhalation.

(iv) Statements reflecting requirements of applicable sections of the National Environmental Protection Act (NEPA), the Superfund Amendments and Reauthorization Act (SARA), and the Occupational Safety and Health Administration's (OSHA) human safety guidance regulations.

(v) Contact address and phone number for reporting adverse reactions or to request a copy of the Materials Safety Data Sheet (MSDS).

[41 FR 38652, Sept. 10, 1976, as amended at 54 FR 18281, Apr. 28, 1989; 61 FR 15704, Apr. 9, 1996]

§ 573.480 Formic acid.

Formic acid may be safely used as a preservative in hay crop silage in an amount not to exceed 2.25 percent of

¹Copies of the current edition may be obtained from the Treasurer, Donald H. James, West Virginia Department of Agriculture, Capital Complex/Guthrie Center, Charleston, WV 25305.

the silage on a dry weight basis or 0.45 percent when direct-cut. The top foot of silage stored should not contain formic acid and silage should not be fed to livestock within 4 weeks of treatment.

§ 573.500 Condensed, extracted glutamic acid fermentation product.

Condensed, extracted glutamic acid fermentation product may be safely used in animal feed under the following conditions:

(a) The additive is a concentrated mixture of the liquor remaining from the extraction of glutamic acid, combined with the cells of *Corynebacterium lilium* used to produce the glutamic acid.

(b) It is used or intended for use as follows:

(1) In poultry feed as a source of protein in an amount not to exceed 5 percent of the total ration.

(2) In cattle feed as a source of protein in an amount not to exceed 10 percent of the feed.

(c) In order to assure safe use, the label and labeling of the additive shall bear, in addition to the other information required by the Act, the following:

(1) The name of the additive.

(2) A statement of the concentration of the additive contained in any mixture.

(3) Adequate directions for use.

§ 573.520 Hemicellulose extract.

Hemicellulose extract may be safely used in animal feed when incorporated therein in accordance with the following conditions:

(a) The additive is produced from the aqueous extract obtained by the treatment of wood with water at elevated temperatures (325 degrees–535 degrees F) and pressure (80 to 900 pounds per square inch) and contains primarily pentose and hexose sugars.

(b) The additive may be used in a liquid or dry state with the liquid product containing not less than 55 percent carbohydrate and the dry product containing not less than 84 percent carbohydrate.

(c) The additive is used as a source of metabolizable energy in animal feed in

accordance with good manufacturing and feeding practices.

[41 FR 38652, Sept. 10, 1976, as amended at 43 FR 11181, Mar. 17, 1978]

§ 573.530 Hydrogenated corn syrup.

(a) *Identity.* The product is produced by hydrogenation of corn syrup over a nickel catalyst.

(b) *Specifications.* The product contains 70 percent hydrogenated corn syrup and a maximum of 0.5 percent reducing sugars.

(c) *Uses.* The product is used as a humectant and plasticizer in preparation of soft-moist dog and cat foods.

(d) *Limitations.* The product is preferably stored in a closed, stainless steel or aluminum container. The level of use of the product shall not exceed 15 percent of the total weight of the pet food formulation.

(e) *Labeling.* The labeling shall bear, in addition to other information required by the Act:

(1) The name of the additive.

(2) Adequate directions for use in accordance with the provisions in paragraph (d) of this section.

[45 FR 22920, Apr. 4, 1980]

§ 573.540 Hydrolyzed leather meal.

(a) *Identity.* Hydrolyzed leather meal is produced from leather scraps that are treated with steam for not less than 33 minutes at a pressure of not less than 125 pounds per square inch.

(b) *Specifications.* The additive shall conform to the following percent-by-weight specifications:

Moisture, not less than 5 percent nor more than 10 percent.

Crude protein, not less than 60 percent.

Crude fat, not less than 5 percent.

Crude fiber, not more than 6 percent.

Chromium, not more than 2.75 percent.

(c) *Use.* It is used or intended for use as a source of protein in swine feeds in an amount not to exceed 1.0 percent by weight of the finished feed.

(d) *Labeling.* The labels and labeling shall bear, in addition to the other information required by the Act:

(1) The name of the additive, hydrolyzed leather meal.

(2) Adequate directions to provide finished feeds complying with paragraph (c) of this section.