

§ 640.22

(b) Plasmapheresis donors shall meet the criteria for suitability prescribed in §640.63, excluding the phrase "other than malaria" in paragraph (c)(9). Informed consent shall be required as prescribed in §640.61.

(c) Plateletpheresis donors shall meet criteria for suitability as described in a license application or a supplement to the product license, and must have the written approval of the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

[40 FR 4304, Jan. 29, 1975, as amended at 49 FR 23834, June 8, 1984; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994]

§640.22 Collection of source material.

(a) Whole blood used as the source of Platelets shall be collected as prescribed in §640.4, except that paragraphs (d)(2) and (h) shall not apply.

(b) If plasmapheresis is used, the procedure for collection shall be prescribed in §§640.62, 640.64 (except paragraph (c)(3)), and 640.65.

(c) If plateletpheresis is used, the procedure for collection shall be as described in a license application or a supplement to a product license, and must have the written approval of the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

(d) The phlebotomy shall be performed by a single uninterrupted venipuncture with minimal damage to, and minimal manipulation of, the donor's tissue.

[40 FR 4304, Jan. 29, 1975, as amended at 45 FR 27927, Apr. 25, 1980; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994]

§640.23 Testing the blood.

(a) Blood from which plasma is separated for the preparation of Platelets shall be tested as prescribed in §§610.40 and 610.45 of this chapter and §640.5 (a), (b), and (c).

(b) The tests shall be performed on a sample of blood collected at the time of collecting the source blood, and such sample container shall be labeled with

21 CFR Ch. I (4-1-98 Edition)

the donor's number before the container is filled.

[40 FR 4304, Jan. 29, 1975, as amended at 50 FR 4139, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988]

§640.24 Processing.

(a) Separation of plasma and platelets and resuspension of the platelets shall be in a closed system. Platelets shall not be pooled during processing.

(b) Immediately after collection, the whole blood or plasma shall be held in storage between 20 to 24 °C, unless it must be transported from the donor clinic to the processing laboratory. During such transport, all reasonable methods shall be used to maintain the temperature as close as possible to a range between 20 and 24 °C until it arrives at the processing laboratory where it shall be held between 20 and 24 °C until the platelets are separated. The platelet concentrate shall be separated within 4 hours after the collection of the unit of whole blood or plasma.

(c) The time and speed of centrifugation must have been demonstrated to produce an unclumped product, without visible hemolysis, that yields a count of not less than 5.5×10^{10} platelets per unit in at least 75 percent of the units tested.

(d) The volume of original plasma used for resuspension of the platelets shall be determined by the maintenance of a pH of not less than 6.0 during the storage period. The pH shall be measured on a sample of platelets which has been stored for the maximum dating period at the selected storage temperature. One of the following storage temperatures shall be used continuously:

(1) 20 to 24 °C.

(2) 1 to 6 °C.

(e) Final containers used for Platelets shall be colorless and transparent to permit visual inspection of the contents; any closure shall maintain a hermetic seal and prevent contamination of the contents. The container material shall not interact with the contents, under the customary conditions of storage and use, in such a manner as to have an adverse effect upon the safety, purity, potency, or efficacy of the product. At the time of filling, the final container shall be marked or identified

by number so as to relate it to the donor.

[40 FR 4304, Jan. 29, 1975, as amended at 42 FR 10983, Feb. 25, 1977; 47 FR 49021, Oct. 29, 1982; 50 FR 4139, Jan. 29, 1985]

§ 640.25 General requirements.

(a) *Storage.* Immediately after re-suspension, Platelets shall be placed in storage at the selected temperature range. If stored at 20 to 24 °C, a continuous gentle agitation of the platelet concentrate shall be maintained throughout the storage period. Agitation is optional if stored at a temperature between 1 and 6 °C.

(b) *Quality control testing.* Each month four units prepared from different donors shall be tested at the end of the storage period as follows:

- (1) Platelet count.
- (2) pH of not less than 6.0 measured at the storage temperature of the unit.
- (3) Measurement of actual plasma volume.
- (4) If the results of the quality control testing indicate that the product does not meet the prescribed requirements, immediate corrective action shall be taken and a record maintained of such action.

(c) *Manufacturing responsibility.* All manufacturing of Platelets shall be performed at the same licensed establishment, except that the quality control testing under paragraph (b) of this section may be performed by a clinical laboratory which meets the standards of the Clinical Laboratories Improvement Act of 1967 (CLIA) (42 U.S.C. 263a) and is qualified to perform platelet counts. Such arrangements must be approved by the Director, Center for Biologics Evaluation and Research, Food and Drug Administration. Such testing shall not be considered as divided manufacturing, as described in § 610.63 of this chapter, provided the following conditions are met:

(1) The results of each test are received within 10 days of the preparation of the platelet concentrate, and are maintained by the establishment licensed for Platelets so that they may be reviewed by an authorized representative of the Food and Drug Administration.

(2) The licensed Platelets manufacturer has obtained a written agreement

that the testing laboratory will permit an authorized representative of the Food and Drug Administration to inspect its testing procedures and facilities during reasonable business hours.

(3) The testing laboratory will participate in any proficiency testing programs undertaken by the Center for Biologics Evaluation and Research, Food and Drug Administration.

[40 FR 4304, Jan. 29, 1975, as amended at 47 FR 49021, Oct. 29, 1982; 49 FR 23834, June 8, 1984; 50 FR 4139, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990]

§ 640.27 Emergency provisions.

The use of the plateletpheresis procedure to obtain a product for a specific recipient may be at variance with §§ 640.21(c) and 640.22(c): *Provided*, That: (a) A licensed physician has determined that the recipient must be transfused with the platelets from a specific donor, and (b) the plateletpheresis procedure is performed under the supervision of a qualified licensed physician who is aware of the health status of the donor and the physician has certified in writing that the donor's health permits plateletpheresis.

[40 FR 53544, Nov. 18, 1975]

Subpart D—Plasma

§ 640.30 Plasma.

(a) *Proper name and definition.* The proper name of this product shall be Plasma. The product is defined as the fluid portion of one unit of human blood intended for intravenous use which in a closed system, has been collected, stabilized against clotting, and separated from the red blood cells.

(b) *Source.* (1) Plasma shall be obtained by separating plasma from blood collected from blood donors or by plasmapheresis.

(2) Plasma may be obtained from a unit of Whole Blood collected by another licensed establishment.

[42 FR 59878, Nov. 22, 1977; 48 FR 13026, Mar. 29, 1983, as amended at 50 FR 4139, Jan. 29, 1985]