

§ 180.367

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting §180.364, see List of CFR Sections Affected in the Finding Aids section of this volume.

§ 180.367 n-Octyl bicycloheptenedicarboximide; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the insecticide *n*-octyl bicycloheptene-dicarboximide, resulting from dermal application, in food commodities as follows:

Commodity	Parts per million
Cattle, fat	0.3
Goats, fat	0.3
Hogs, fat	0.3
Horses, fat	0.3
Milk, fat	0.3
Sheep, fat	0.3

(2) *N*-octylbicycloheptene dicarboximide may be safely used in accordance with the following prescribed conditions:

(i) It is used in combination with piperonyl butoxide and pyrethrins for insect control in food-processing and food-storage areas, provided that the food is removed or covered prior to such use.

(ii) Residues in food resulting from the use described in paragraph (a)(2)(i) of this section shall not exceed 10 parts per million of *N*-octylbicycloheptene dicarboximide, 10 parts per million of piperonyl butoxide, and 1 part per million of pyrethrins.

(iii) To assure safe use of the pesticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency and it shall be used in accordance with such label and labeling.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[65 FR 33713, May 24, 2000]

§ 180.368 Metolachlor; tolerances for residues.

(a) *General.* Tolerances are established for the combined residues (free and bound) of the herbicide metolachlor [2-chloro-*N*-(2-ethyl-6-methylphenyl)-*N*-(2-methoxy-1-

40 CFR Ch. I (7-1-00 Edition)

methylethyl)acetamide] and its metabolites, determined as the derivatives, 2-[(2-ethyl-6-methylphenyl)amino]-1-propanol and 4-(2-ethyl-6-methylphenyl)-2-hydroxy-5-methyl-3-morpholinone, each expressed as the parent compound, in or on the raw agricultural commodities.

Commodity	Parts per million
Almond hulls	0.3
Barley, fodder	0.5
Barley, grain	0.1
Buckwheat, grain	0.1
Cabbage	1.0
Cattle, fat	0.02
Cattle, kidney	0.2
Cattle, liver	0.05
Cattle, meat	0.02
Cattle, mbyp (except kidney and liver)	0.02
Celery	0.1
Corn, fresh (inc. sweet K=CWHR)	0.1
Corn, forage and fodder	8.0
Corn, grain	0.1
Cottonseed	0.1
Eggs	0.02
Goats, fat	0.02
Goats, kidney	0.2
Goats, liver	0.05
Goats, meat	0.02
Goats, mbyp (except kidney and liver)	0.02
Hogs, fat	0.02
Hogs, kidney	0.2
Hogs, liver	0.05
Hogs, meat	0.02
Hogs, mbyp (except kidney and liver)	0.02
Horses, fat	0.02
Horses, kidney	0.2
Horses, liver	0.05
Horses, meat	0.02
Horses, mbyp (except kidney and liver)	0.02
Legume vegetables group foliage (except soybean forage and soybean hay)	15.0
Milk	0.02
Millet, fodder	0.5
Millet, forage	0.5
Millet, grain	0.1
Milo, fodder	0.5
Milo, forage	0.5
Milo, grain	0.1
Nongrass animal feeds (forage, fodder, straw, and hay) group	3.0
Oats, fodder	0.5
Oats, forage	0.5
Oats, grain	0.1
Peanuts	0.5
Peanut, forage and hay	30.0
Peppers, bell	0.1
Potatoes	0.2
Poultry, fat	0.02
Poultry, liver	0.05
Poultry, meat	0.02
Poultry, mbyp (except liver)	0.02
Rice, fodder	0.5
Rice, forage	0.5
Rice, grain	0.1
Rye, fodder	0.5
Rye, forage	0.5
Rye, grain	0.1
Safflower seed	0.1
Seed and pod vegetables (except soybeans)	0.3
Sheep, fat	0.02