

§ 180.409

40 CFR Ch. I (7-1-00 Edition)

| Commodity  | Part per million |
|--|------------------|
| Barley, grain .....  | 0.2              |
| Barley, fodder .....   | 2.0              |
| Barley, milling fractions .....                                    | 1.0              |
| Barley, straw .....  | 2.0              |
| Cereal grains group (except wheat, barley, and oats), fodder ..... | 1.0              |
| Cereal grains group (except wheat, barley, and oats), forage ..... | 1.0              |
| Cereal grains group (except wheat, barley, and oats), straw .....  | 1.0              |
| Oat, fodder .....  | 2.0              |
| Oat, forage .....  | 2.0              |
| Oat, grain .....   | 0.2              |
| Oat milling fractions .....  | 1.0              |
| Oat, straw .....   | 2.0              |
| Wheat, fodder .....  | 2.0              |
| Wheat, forage .....  | 2.0              |
| Wheat, grain .....   | 0.2              |
| Wheat, milling fractions .....                                     | 1.0              |
| Wheat, straw .....   | 2.0              |

[65 FR 33700, May 24, 2000]

§ 180.409 Pirimiphos-methyl; tolerances for residues.

(a) *General.* (1) Tolerances are established for the combined residues of the insecticide pirimiphos-methyl, *O*-[2-diethylamino-6-methyl-4-pyrimidinyl] *O,O*-dimethyl phosphorothioate, the metabolite *O*-(2-ethylamino-6-methylpyrimidin-4-yl) *O,O*-dimethyl phosphorothioate and, in free and conjugated form, the metabolites 2-diethylamino-6-methyl-pyrimidin-4-ol, 2-ethylamino-6-methyl-pyrimidin-4-ol, and 2-amino-6-methyl-pyrimidin-4-ol in or on the following food commodities:

| Commodity                                   | Parts per million |
|---|-------------------|
| Corn .....                                  | 8.0               |
| Cattle fat .....                            | 0.2               |
| Cattle, kidney and liver .....              | 2.0               |
| Cattle, mbyop .....                         | 0.2               |
| Cattle, meat .....                          | 0.2               |
| Eggs .....                                  | 0.5               |
| Goats, fat .....                            | 0.2               |
| Goats, kidney and liver .....               | 2.0               |
| Goats, mbyop .....                          | 0.2               |
| Goats, meat .....                           | 0.2               |
| Hogs, fat .....                             | 0.2               |
| Hogs, kidney and liver .....                | 2.0               |
| Hogs, mbyop .....                           | 0.2               |
| Hogs, meat .....                            | 0.2               |
| Horses, fat .....                           | 0.2               |
| Horses, kidney and liver .....              | 2.0               |
| Horses, mbyop .....                         | 0.2               |
| Horses, meat .....                          | 0.2               |
| Kiwifruit .....                             | 5.0               |
| Milk, fat (0.1 ppm (N) in whole milk) ..... | 3.0               |
| Poultry, fat .....                          | 0.2               |
| Poultry, mbyop .....                        | 2.0               |
| Poultry, meat .....                         | 2.0               |
| Sheep, fat .....                            | 0.2               |
| Sheep, kidney and liver .....               | 2.0               |
| Sheep, mbyop .....                          | 0.2               |

| Commodity            | Parts per million |
|----------------------|-------------------|
| Sheep, meat .....    | 0.2               |
| Sorghum, grain ..... | 8.0               |

(2) Tolerances are established for the combined residues of the insecticide pirimiphos-methyl (*O*-[2-diethylamino-6-methyl-4-pyrimidinyl] *O,O*-dimethyl phosphorothioate) and its metabolite *O*-(2-ethylamino-6-methyl-pyrimidin-4-yl) *O,O*-dimethyl phosphorothioate and, in free and conjugated forms, the metabolites 2-diethylamino-6-methyl-pyrimidin-4-ol, 2-ethylamino-6-methyl-pyrimidin-4-ol, and 2-amino-6-methyl-pyrimidin-4-ol in or on the following food commodities when present therein as a result of application to stored grains:

| Food   | Parts per million |
|--|-------------------|
| Corn milling fractions (except flour) .....    | 40                |
| Corn oil .....                                 | 88                |
| Sorghum milling fractions (except flour) ..... | 40                |

(3) A tolerance of 8.0 parts per million is established for residues of the insecticide pirimiphos-methyl (*O*-[2-diethylamino-6-methyl-4-pyrimidinyl] *O,O*-dimethyl phosphorothioate) and its metabolite *O*-(2-ethylamino-6-methyl-pyrimidin-4-yl) *O,O*-dimethyl phosphorothioate and, in free and conjugated forms, the metabolites 2-diethylamino-6-methyl-pyrimidin-4-ol, 2-ethylamino-6-methyl-pyrimidin-4-ol, and 2-amino-6-methyl-pyrimidin-4-ol in or on the processed commodity wheat flour as a result of application to stored wheat grain. There are no U.S. registrations for use of pirimiphos-methyl on wheat, as of June 12, 1990.

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[65 FR 33714, May 24, 2000]

§ 180.410 Triadimefon; tolerances for residues.

(a) *General.* Tolerances are established for the combined residues of the fungicide 1-(4-chlorophenoxy)-3,3-dimethyl-1*H*-1,2,4-triazol-1-yl)-2-butanone and its metabolites containing