

§ 147.22 Hatching egg sanitation.

Hatching eggs should be collected from the nests at frequent intervals and, to aid in the prevention of contamination with disease-causing organisms, the following practices should be observed:

(a) Cleaned and disinfected containers, such as egg flats, should be used in collecting the nest eggs for hatching. Egg handlers should thoroughly wash their hands with soap and water prior to and after egg collection. Clean outer garments should be worn.

(b) Dirty eggs should not be used for hatching purposes and should be collected in a separate container from the nest eggs. Slightly soiled nest eggs may be gently dry cleaned by hand.

(c) Hatching eggs should be stored in a designated egg room under conditions that will minimize egg sweating. The egg room walls, ceiling, floor, door, heater, and humidifier should be cleaned and disinfected after every egg pickup. Cleaning and disinfection procedures should be as outlined in § 147.24.

(d) The egg processing area should be cleaned and disinfected daily.

(e) Effective rodent and insect control programs should be implemented.

(f) The egg processing building or area should be designed, located, and constructed of such materials as to assure that proper egg sanitation procedures can be carried out, and that the building itself can be easily, effectively, and routinely sanitized.

(g) All vehicles used for transporting eggs or chicks/poults should be cleaned and disinfected after use. Cleaning and disinfection procedures should be as outlined in § 147.24.

[67 FR 8474, Feb. 25, 2002]

§ 147.23 Hatchery sanitation.

An effective program for the prevention and control of *Salmonella* and other infections should include the following measures:

(a) An effective hatchery sanitation program should be designed and implemented.

(b) The hatchery building should be arranged so that separate rooms are provided for each of the four operations: Egg receiving, incubation and hatching, chick/poult processing, and

egg tray and hatching basket washing. Traffic and airflow patterns in the hatchery should be from clean areas to dirty areas (*i.e.*, from egg room to chick/poult processing rooms) and should avoid tracking from dirty areas back into clean areas.

(c) The hatchery rooms, and tables, racks, and other equipment in them should be thoroughly cleaned and disinfected frequently. All hatchery wastes and offal should be burned or otherwise properly disposed of, and the containers used to remove such materials should be cleaned and sanitized after each use.

(d) The hatching compartments of incubators, including the hatching trays, should be thoroughly cleaned and disinfected after each hatch.

(e) Only clean eggs should be used for hatching purposes.

(f) Only new or cleaned and disinfected egg cases should be used for transportation of hatching eggs. Soiled egg case fillers should be destroyed.

(g) Day-old chicks, poults, or other newly hatched poultry should be distributed in clean, new boxes and new chick papers. All crates and vehicles used for transporting birds should be cleaned and disinfected after each use.

[67 FR 8474, Feb. 25, 2002]

§ 147.24 Cleaning and disinfecting.

The following procedures are recommended:

(a) In the poultry houses:

(1) Remove all live "escaped" and dead birds from the building. Blow dust from equipment and other exposed surfaces. Empty the residual feed from the feed system and feed pans and remove it from the building. Disassemble feeding equipment and dump and scrape as needed to remove any and all feed cake and residue. Clean up spilled feed around the tank and clean out the tank. Rinse down and wash out the inside of the feed tank to decontaminate the surfaces and allow to dry.

(2) Remove all litter and droppings to an isolated area where there is no opportunity for dissemination of any infectious disease organisms that may be present. Housing where poultry infected with a mycoplasmal disease were kept should remain closed for 7 days before removal of the litter.